

Agrisera

This product is for research use only (not for diagnostic or therapeutic use)

contact: support@agrisera.com

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | www.agrisera.com

Product no **AS16 3169** **H3T3pK4me2 | Histone H3 (dimethylated Lys4, p Thr3)**


Product information

Immunogen	KLH-conjugated synthetic peptide
Host	Rabbit
Clonality	Polyclonal
Purity	Affinity purified serum
Format	Liquid
Quantity	50 µg
Storage	Store lyophilized/reconstituted at -20°C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please, remember to spin tubes briefly prior to opening them to avoid any losses that might occur from lyophilized material adhering to the cap or sides of the tubes.
Additional information	This antibody preparation is provided in 20 mM Potassium Phosphate pH 7.2, 150 mM NaCl, 0,01% sodium azide and 30% glycerol.


Application information

Recommended dilution	2-5 µg/million cells (ChiP), 1 : 1000 (Dot), 1 : 50 (IF), 1 : 50 (IHC), 1 : 500 (WB)
Expected apparent MW	15 kDa
Confirmed reactivity	Human
Predicted reactivity	<i>Caenorhabditis elegans</i> , <i>Drosophila melanogaster</i> , Mouse, Plant, Rat, <i>Xenopus sp.</i>
Not reactive in	No confirmed exceptions from predicted reactivity are currently known.

application example



Chromatin Immunoprecipitation using H3T3pK4me2 antibodies. Chromatin from one million formaldehyde cross-linked HeLa cells was used with 2 µg of H3T3pK4me2 and 20 µl of magnetic beads per immunoprecipitation. A no antibody (No Ab) control was also used. Immunoprecipitated DNA was quantified using quantitative real-time PCR and normalized to the input chromatin.




Immunofluorescence using anti-H3T3pK4me2 antibodies. Tissue: HeLa cells. Fixation: 0.5% PFA. Primary antibody was used at a 1:50 dilution for 1 h at RT. Secondary antibody: Dylight@ 488 secondary antibody at 1:10 000 for 45 min at RT. Localization: Histone H3T3pK4me2 is nuclear and chromosomal. Staining: H3T3pK4me2 is expressed in green while the nuclei and alpha-tubulin were coexpressed with DAPI (blue) and Dylight@ 550 (red).

Agrisera

This product is **for research use only** (not for diagnostic or therapeutic use)

contact: support@agrisera.com

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | www.agrisera.com



Western Blot using anti-H3T3pK4me2 antibodies. 30 μ g of NIH-3T3 histone extracts. Primary antibody used at a 1:500 dilution overnight at 4 °C. Secondary antibody: IRDye800™ rabbit secondary antibody at 1:10 000 for 45 min at RT.