

This product is **for research use only** (not for diagnostic or therapeutic use)

contact: support@agrisera.com

Agrisera AB | Box 57 | SE-91112 Vännäs | Sweden | +46 (0)935 33 000 | www.agrisera.com

Product no **AS07 251**

Anti-FtsH10 | ATP-dependent zinc metalloprotease FtsH10 (mitochondrial)


Product information

Immunogen	KLH-conjugated peptide located near C-terminus chosen from sequence of <i>Arabidopsis thaliana</i> FtsH10 <u>Q8VZI8</u> . <u>At1g07510</u>
Host	Rabbit
Clonality	Polyclonal
Purity	Serum
Format	Lyophilized
Quantity	200 µl
Reconstitution	For reconstitution add 200 µl of sterile water
Storage	Store lyophilized/reconstituted at -20°C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please remember to spin the tubes briefly prior to opening them to avoid any losses that might occur from material adhering to the cap or sides of the tube.
Additional information	Blue-native (2D BN/SDS-PAGE) methodology has been described in Piechota et al, 2010

Application information

Recommended dilution	1 : 1000 (WB)
Expected apparent MW	84 kDa
Confirmed reactivity	<i>Arabidopsis thaliana</i>
Predicted reactivity	<i>Arabidopsis thaliana</i>
Not reactive in	No confirmed exceptions from predicted reactivity are currently known
Selected references	<p>Kolodziejczak et al. (2018). m-AAA Complexes Are Not Crucial for the Survival of <i>Arabidopsis</i> Under Optimal Growth Conditions Despite Their Importance for Mitochondrial Translation. <i>Plant Cell Physiol.</i> 2018 May 1;59(5):1006-1016. doi: 10.1093/pcp/pcy041.</p> <p>Piechota et al. (2015). Unraveling the functions of type II-prohibitins in <i>Arabidopsis</i> mitochondria. <i>Plant Mol Biol.</i> 2015 Apr 21.</p> <p>Kwasniak et al. (2013). Silencing of the Nuclear RPS10 Gene Encoding Mitochondrial Ribosomal Protein Alters Translation in <i>Arabidopsis</i> Mitochondria. <i>Plant Cell</i>, May 30.</p> <p>Quesada et al. (2011). <i>Arabidopsis</i> RUGOSA2 encodes an mTERF family member required for mitochondrion, chloroplast and leaf development. <i>Plant J.</i></p>

Application example



Mitochondrial preparation from *Arabidopsis thaliana* mitochondria was separated on 10% acrilamide gel and electrophoresis prepared according to Schägger and von Jagow (Anl. Biochem., 1987, 166:368-379). After running the gel, proteins were transferred to nitrocellulose membrane using wet transfer (0.22% CAPS, pH 11). Transfer was checked by Ponceau S staining. Blot was destained by several quick washings in distilled water and 1 washing in 1X TBS (10 mM T pH 7.5, 150 mM NaCl) (10-15 min.). Blot was blocked by 1.5 hour in 5% milk in TBST (1X TBS, 0,1 20) After blocking blot was washed quickly twice in TBST and incubated 2 hours with primary antibody (dilution 1: 1000 TBST (dilution 1:1000)). Washing: two quick washings in TBST and 3 x 10 min. washings in TBST. Then blot was incubated 45-60 min. with a secondary anti-rabbit antibodies conjugated to peroxidase (dilution 1:10000) in TBST. Washing: as above. After washing blot was incubated 1-2 min. in ECL solution and exposed to Kodak autoradiography film. Exposure time was 15-60 seconds.



This product is **for research use only** (not for diagnostic or therapeutic use)

contact: support@agrisera.com

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | www.agrisera.com

Mitochondria were isolated as described by Urantowka et al. (Plant Mol Biol, 2005, 59:239-52). Mitochondrial pellets were suspended in 1X Laemmli buffer (5% beta-mercaptoethanol, 3.7% glycerol, 1.1% SDS, 23 mM Tris-HCl pH 6.8, 0.01% bromophenol blue), heated (95 °C, 5 min.) and centrifuged (13000 rpm, 1 min.).

Courtesy Dr. J. Piechota, University of Wroclaw, Poland