

# Agrisera

This product is for research use only (not for diagnostic or therapeutic use)

contact: [support@agrisera.com](mailto:support@agrisera.com)

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | [www.agrisera.com](http://www.agrisera.com)

Product no **AS10 673**  
**AGO9 | Argonaute 9**


## Product information

<b>Immunogen</b>	KLH-conjugated synthetic peptide derived from <i>Arabidopsis thaliana</i> AGO9 protein sequence UniProt: <a href="#">Q84YI4</a> , TAIR: <a href="#">At5g21150</a> .
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Purity</b>	Affinity purified serum in PBS, pH 7.4
<b>Format</b>	Lyophilized
<b>Quantity</b>	100 µg
<b>Reconstitution</b>	For reconstitution add 100 µl of sterile water.
<b>Storage</b>	Store lyophilized/reconstituted at -20 °C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please, remember to spin tubes briefly prior to opening them to avoid any losses that might occur from lyophilized material adhering to the cap or sides of the tubes.

## Application information

<b>Recommended dilution</b>	5 µg of antibody per 1 gram of a fresh tissue (IP), 1 : 10 000 (WB)
<b>Expected   apparent MW</b>	101 kDa
<b>Confirmed reactivity</b>	<i>Arabidopsis thaliana</i>
<b>Predicted reactivity</b>	<i>Arabidopsis thaliana</i>
<b>Not reactive in</b>	No confirmed exceptions from predicted reactivity are currently known.
<b>Additional information</b>	AGO expression may be cell/tissue specific and using floral tissue is recommended where most of the AGOs are expressed the highest. Seedlings can be used as a negative control. Use of proteasome inhibitors as MG132 can help to stabilize AGO proteins during extraction procedure. For high resolution images, please visit the specific product page at <a href="http://www.agrisera.com">www.agrisera.com</a>
<b>Selected references</b>	<a href="#">Sprunck et al. (2019)</a> . Elucidating small RNA pathways in <i>Arabidopsis thaliana</i> egg cells. <a href="http://dx.doi.org/10.1101/525956">http://dx.doi.org/10.1101/525956</a> <a href="#">Su et al. (2017)</a> . The THO Complex Non-Cell-Autonomously Represses Female Germline Specification through the TAS3-ARF3 Module. <i>Curr Biol.</i> 2017 Jun 5;27(11):1597-1609.e2. doi: 10.1016/j.cub.2017.05.021. <a href="#">Havecker et al. (2010)</a> The RNA-directed DNA methylation <i>Arabidopsis</i> Argonautes functionally diverge based on expression and interaction with target loci. <i>Plant Cell</i> 22(2): 321-34.

## Application example



80 µg of *Arabidopsis thaliana* soluble total cell extract (extracted in 20 mM Tris pH7.5, 5mM MgCl<sub>2</sub>, 2.5mM DTT, 300mM NaCl, 0.1% NP-40, 1% proteaseinhibitor) was separated on 6% SDS-PAGE and blotted 1h to **nitrocellulose**. Filters were blocked 1h with 5% low-fat milk powder in

# Agrisera

This product is **for research use only** (not for diagnostic or therapeutic use)

contact: [support@agrisera.com](mailto:support@agrisera.com)

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | [www.agrisera.com](http://www.agrisera.com)

TBS-TT (0.25% TWEEN20; 0.1% Triton-X) and probed with anti-AGO9 antibody (1:10 000, 1h) and secondary anti-rabbit antibody (HRP conjugated, Agrisera [AS09\\_602](#)) (1:15 000, 1 h) in TBS-TT containing 5% low fat milk powder. Antibody incubations were followed by washings in TBS-TT. All steps were performed at RT with agitation. Blots were developed for 5 min with chemiluminescent detection reagent according the manufacturer's instructions. Exposure time was 30 seconds.

Courtesy Dr. Ericka Havecker, University of Cambridge, UK