Agrisera
Part of Thermo Fisher Scientific

This product is **for research use only** (not for diagnostic or therapeutic use) **contact: support@agrisera.com**

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | www.agrisera.com

Product no AS12 2608

Anti-TOR | Target of rapamycin

Product information

Immunogen	KLH-conjugated synthetic peptide derived from *Arabidopsis thaliana* TOR protein sequence, UniProt: Q9FR53, TAIR: AT1G50030
Host	Rabbit
Clonality	Polyclonal
Purity	Immunogen affinity purified serum in PBS pH 7.4.
Format	Lyophilized
Quantity	50 µg
Reconstitution	For reconstitution add 50 µl of sterile water
Storage	Store lyophilized/reconstituted at -20°C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please remember to spin the tubes briefly prior to opening them to avoid any losses that might occur from material adhering to the cap or sides of the tube.

Application information

Recommended dilution	1 : 5000 (WB)
Expected | apparent MW	279 kDa
Confirmed reactivity	*Arabidopsis thaliana, Zea mays*
Predicted reactivity	*Cucumis sativus, Glycne max, Nicotiana tabacum, Oryza sativa, Phaseolus vulgaris, Physcomitrium patens, Populus trichocarpa, Setaria italica, Setaria viridis, Solanum tuberosum, Sorghum bicolor, Vitis vinifera* Species of your interest not listed? Contact us
Not reactive in	*Chlamydomonas reinhardtii*
Additional information	TOR is a subjected to post-translational modifications (ubiquitination and phosphorylation), which can change the migration pattern in SDS gel
Selected references	Liu et al. (2025). Guard cell and whole plant expression of AtTOR improves performance under drought and enhances water use efficiency. J Biol Chem. 2025 Jun;301(6):110220. doi: 10.1016/j.jbc.2025.110220. Garcia et al. (2017). Maize defective kernel mutant generated by insertion of a Ds element in a gene encoding a highly conserved TTI2 cochaperone. Proc Natl Acad Sci U S A. 2017 May 16;114(20):5165-5170. doi: 10.1073/pnas.1703498114.

application information



25 mg total protein extract (35S:Raptor-ProtG) was incubated with IgG magnetic beads for 50 minutes. After washing, complexes were eluted, concentrated by speedvac and pellet was dissolved in 30uL 1x sample buffer. For SDS-PAGE 10uL of the eluate was loaded on gel. 20, 40, and 60 µg total protein extract from *Arabidopsis thaliana* cell suspension culture (Landsberg erecta), or AtTOR protein enriched by immunoprecipitation using AtRAPTOR1B was separated on 4-15 % gradient TGX (Biorad) SDS-PAGE and blotted 15 min. to PVDF using the Trans-Blot Turbo system (setting high MW). As negative control, an unrelated immunoprecipitated sample was loaded. Blots were blocked overnight with 1% PVP40 in TBS-T at 4°C with agitation. Blot was incubated with the primary anti-TOR antibody at a dilution of 1: 1 000 in blocking buffer for 4h at room temperature (RT) with agitation. The antibody solution was decanted and the blot was rinsed briefly once, then washed once for 15 min and 3 times for 5 min in TBS-T at RT with agitation. Blot was incubated in secondary antibody (anti-rabbit IgG horse radish peroxidase conjugated, from Agrisera AS09 602) diluted to 1: 20 000 in blocking buffer for 1h at RT with agitation. The blot was washed as above and developed for 1 min with ECL according to the manufacturer's instructions. Exposure time was 10 seconds.

TOR protein (red arrow) detected after enrichment by immunoprecipitation with IgG beads and Raptor-ProtG fusion, while TOR band is

Agrisera
Part of Thermo Fisher Scientific

This product is **for research use only** (not for diagnostic or therapeutic use) **contact: support@agrisera.com**

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | www.agrisera.com

absent in control IP with unrelated ProtG-tagged bait.