

This product is **for research use only** (not for diagnostic or therapeutic use)

contact: support@agrisera.com

Agrisera AB | Box 57 | SE-91112 Vännäs | Sweden | +46 (0)935 33 000 | www.agrisera.com

Product no AS12 2119

Anti-14-3-3 GRF | General regulatory element

Product information

Immunogen	Conserved surface-exposed peptide conjugated to KLH. This sequence is repeated in all 12 "canonical" members of the <i>Arabidopsis thaliana</i> GRF/14-3-3 family. GRF1 GF14 chi UniProt: F4JJ94, TAIR: AT4G09000, GRF2 GF14 omega, UniProt: Q01525, TAIR: AT1G78300, GRF3 GF14 psi UniProt: F4KBI7, TAIR: AT5G38480, GRF4 GF14 phi UniProt: F4HWQ5, TAIR: AT1G35160, GRF5 GF14 upsilon UniProt: P42645, TAIR: AT5G16050, GRF6 GF14 lambda UniProt: P48349#P48349-2, TAIR: AT5G10450, GRF7 GF14 nu UniProt: Q96300, TAIR: AT3G02520, GRF8 GF14 kappa UniProt: F4KHY7, TAIR: AT5G65430, GRF9 GF14 mu UniProt: F4IP55, TAIR: AT2G42590, GRF10 GF14 epsilon UniProt: F4I1C1, TAIR: AT1G22300, GRF11 GF14 omicron UniProt: Q9S9Z8, TAIR: AT1G34760, GRF12 GF14 iota UniProt: Q9C5W6 TAIR: AT1G26480.
Host	Rabbit
Clonality	Polyclonal
Purity	Serum
Format	Lyophilized
Quantity	50 µl
Reconstitution	For reconstitution add 50 µl of sterile water
Storage	Store lyophilized/reconstituted at -20°C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please remember to spin the tubes briefly prior to opening them to avoid any losses that might occur from material adhering to the cap or sides of the tube.
Additional information	<p>This antibody is recognizing recombinant GRF of <i>Lilium longiflorum</i> Lili1433_0 accession: AF191746, Lili1433_2, accession: EF397608 and recombinant GRF1,2,3,5 and 6 of <i>Arabidopsis thaliana</i>: GRF1 14-3-3 chi (At4g09000.1), GRF2 14-3-3 omega (At1g78300.1), GRF3 14-3-3 psi (At5g38480.1), GRF5 14-3-3 upsilon (At5g16050.1), GRF6 14-3-3 lambda (At5g10450.2), GRF8 14-3-3 kappa (At5g65430.1), GRF11 14-3-3 omicron (At1g34760.1).</p> <p>There is also very weak reaction to <i>Physcomitrella patens</i> Pp14-3-3 Pp1s 73_133V6 (closest homolog to AtGRF6) and <i>Chlamydomonas reinhardtii</i> Cr 14-3-3 Cre 12.g559250 (closest homolog to AtGRF6).</p>

Application information


Recommended dilution	1: 2000 (WB)
Expected apparent MW	20-28 kDa (depending upon an isoform)
Confirmed reactivity	<i>Aesculus hippocastanum</i> , <i>Arabidopsis halleri</i> , <i>Arabidopsis thaliana</i> , <i>Lilium longiflorum</i> , <i>Mesembryanthemum crystallinum</i> , <i>Neochloris oloabundans</i> (algae), <i>Oryza sativa</i> , <i>Picrorhiza kurroa</i> , <i>Phaseolus vulgaris</i> , <i>Panax notoginseng</i> , <i>Zea mays</i> , <i>Vicia faba</i>
Predicted reactivity	<i>Brassica</i> sp., <i>Chlamydomonas reinhardtii</i> , <i>Glycine max</i> , <i>Gossypium hirsutum</i> , <i>Nicotiana tabacum</i> , <i>Physcomitrium patens</i> , <i>Pisum sativum</i> , <i>Solanum tuberosum</i> , <i>Triticum aestivum</i>
	Species of your interest not listed? Contact us
Not reactive in	This antibody does not bind to 14-3-3-like protein D of <i>Nicotiana tabacum</i> .
Additional information	This product can be sold containing proclin if requested
Selected references	<p>Guo et al. (2022) Acetylproteomics analyses reveal critical features of lysine-?-acetylation in <i>Arabidopsis</i> and a role of 14-3-3 protein acetylation in alkaline response. <i>Stress Biology</i> 2, 1 (2022). https://doi.org/10.1007/s44154-021-00024-z. (immunoprecipitation)</p> <p>Kumari et al. (2021) In-depth assembly of organ and development dissected <i>Picrorhiza kurroa</i> proteome map using mass spectrometry. <i>BMC Plant Biol.</i> 2021 Dec 22;21(1):604. doi: 10.1186/s12870-021-03394-8. PMID: 34937558; PMCID: PMC8693493.</p> <p>Dongxu et al. (2020). Magnesium reduces cadmium accumulation by decreasing the nitrate reductase-mediated nitric oxide production in <i>Panax notoginseng</i> roots. <i>Journal of Plant Physiology</i>. Available online 7 February 2020, 153131</p> <p>Gupta and Shaw (2020). Biochemical and molecular characterisations of salt tolerance components in rice varieties tolerant and sensitive to NaCl: the relevance of Na⁺ exclusion in salt tolerance in the species . <i>Funct Plant Biol.</i> 2020 Jul 30.doi: 10.1071/FP20089</p> <p>Perl-Obermeyer et al. (2018). Dissecting the subcellular membrane proteome reveals enrichment of H⁺ (co-)transporters and vesicle trafficking proteins in acidic zones of Chara internodal cells. <i>PLoS One.</i> 2018 Aug 29;13(8):e0201480. doi: 10.1371/journal.pone.0201480.</p>

This product is **for research use only** (not for diagnostic or therapeutic use)

contact: support@agrisera.com

Agrisera AB | Box 57 | SE-911121 Vännäs | Sweden | +46 (0)935 33 000 | www.agrisera.com

Application example



Left panel: 150 ng of recombinant *Arabidopsis thaliana*, *Physcomitrella patens* and *Chlamydomonas reinhardtii* GRFs (see more details below) were separated on 12 % SDS-PAGE and blotted 1h to PVDF by semi dry blotting (0.8 mA/cm² for 60 min). Blots were blocked with 2% skimmed milk powder dissolved in TBST (0.1 % Tween 20) for 1h at room temperature (RT) with agitation. Blot was incubated in the primary antibody at a dilution of 1: 1 000 in blocking solution and incubated for 1h at RT with agitation. The antibody solution was decanted and the blot was rinsed briefly twice, then washed once for 15 min and 3 times for 5 min in TBS-T at RT with agitation. Blot was incubated in secondary antibody (anti-rabbit IgG horse radish peroxidase conjugated, from Agrisera AS09 602) diluted to 1:20 000 in for 1h at RT with agitation. The blot was washed as above and developed for 5 min with ECL Prime western blotting detection reagent according to the manufacturer's instructions. Exposure time was 10 minutes.

Right panel: 50 µg of a total cell extract of *Arabidopsis thaliana* wilde type. Conditions as above, exposure time 30 seconds.

GRF protein designations: At 14-3-3 lambda (GRF6), At 14-3-3 chi (GRF1), At14-3-3 psi (GRF3), At14-3-3 omega (GRF2), At14-3-3 upsilon (GRF5), At14-3-3 omicron (GRF11), At14-3-3 kappa (GRF8), Pp 14-3-3 Pp1s 73_133V6 (closest homolog to At GRF6), Cr 14-3-3 Cre 12.g559250 (closest homolog to AtGRF6). Note that all recombinant GRFs were detected by this antibody, At14-3-3 kappa (GRF8) to a lesser extent.

Courtesy of Dr. Bernhard Wurzinger, University of Vienna, Austria