


Product no **AS11 1783****Anti-ACD1 | Accelerated cell death 1****Product information**


<b>Immunogen</b>	Recombinant PaO from <i>Arabidopsis thaliana</i> <a href="#">Q9FYC2</a> , <a href="#">At3g44880</a>
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Purity</b>	Serum
<b>Format</b>	Lyophilized
<b>Quantity</b>	50 µl
<b>Reconstitution</b>	For reconstitution add 50 µl of sterile water
<b>Storage</b>	Store lyophilized/reconstituted at -20°C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please remember to spin the tubes briefly prior to opening them to avoid any losses that might occur from material adhering to the cap or sides of the tube.
<b>Additional information</b>	The protein level is moderately induced during dark-induced senescence

**Application information**

<b>Recommended dilution</b>	1 : 5000 (WB)
<b>Expected   apparent MW</b>	61   54 kDa
<b>Confirmed reactivity</b>	<i>Arabidopsis thaliana</i>
<b>Predicted reactivity</b>	<i>Brassica napus</i> , <i>Solanum lycopersicum</i> , <i>Nicotiana tabacum</i> Species of your interest not listed? <a href="#">Contact us</a>
<b>Not reactive in</b>	<i>Pinus strobus</i>
<b>Additional information</b>	This antibody works on total cell extracts and can be used as a senescence marker. Predicted size of Acd1 precursor protein is about 61 kD including the transit peptide, but it must be processed to a smaller size. Using fresh extracts is recommended to decrease possible cross-reaction with Rubisco.
<b>Selected references</b>	<a href="#">Fukura</a> et al. (2021). Enrichment of chlorophyll catabolic enzymes in grana margins and their cooperation in catabolic reactions. <i>J Plant Physiol.</i> 2021 Nov;266:153535. doi: 10.1016/j.jplph.2021.153535. Epub 2021 Sep 25. PMID: 34607178. <a href="#">Kim</a> et al. (2013). Mutation of the Arabidopsis NAC016 Transcription Factor Delays Leaf Senescence. <i>Plant Cell Physiol.</i> Aug 21. <a href="#">Nagane</a> et al. (2010). Involvement of AtNAP1 in the regulation of chlorophyll degradation in Arabidopsis thaliana. <i>Planta</i> (4):939-949. <a href="#">Hirashima</a> et al. (2009). Light-independent cell death induced by accumulation of pheophorbide a in Arabidopsis thaliana. <i>Plant Cell Physiol.</i> (4):719-729.

**application example**

*Arabidopsis thaliana* wild ecotype Columbia was grown for four weeks under continuous illumination and then transferred to complete darkness for five days. Several leaves were harvested from the plants before they were transferred to darkness (0 d) or after they were kept for five days (5 d). Protein was extracted with the SDS extraction solution containing 50 mM Tris (pH 6.8), 10% (w/v) glycerol, 2% (w/v) SDS and 6% (v/v) 2-mercaptoethanol. Protein extract equivalent to 1 mg leaf material was loaded and separated on 14% SDS-PAGE and blotted 1h to PVDF. Blots were blocked with PBS-T containing 1.5% skim milk for 1h at room temperature (RT) with agitation. Blot was incubated in the primary antibody at a dilution of 1: 10 000 for 1h at RT with agitation. The antibody solution was decanted and the blot was rinsed briefly twice, then washed once for 15 min and 3 times for 5 min in PBS-T at RT with agitation. Blot was incubated in secondary antibody (anti-rabbit IgG horse radish peroxidase conjugated, from GE Healthcare) diluted to 1:20 000 in for 1h at RT with agitation. The blot was washed as above and developed for 1 min with ECLplus according to the manufacturers instructions. Exposure time was 5 min.



*Arabidopsis thaliana* wild ecotype Columbia was grown for four weeks under continuous illumination. Several young (**1**), mature (**2**) and senescing (**3**) leaves were harvested from the plants. Protein was extracted with the SDS extraction solution containing 50 mM Tris (pH 6.8), 10% (w/v) glycerol, 2% (w/v) SDS and 6% (v/v) 2-mercaptoethanol. Protein extract equivalent to 1 mg leaf material was loaded and separated on 14% SDS-PAGE and blotted 1h to PVDF. Blots were blocked with PBS-T containing 1.5% skim milk for 1h at room temperature (RT) with agitation. Blot was incubated in the primary antibody at a dilution of 1:30 000 for 1h at RT with agitation as indicated in the figure. The antibody solution was decanted and the blot was rinsed briefly twice, then washed once for 15 min and 3 times for 5 min in PBS-T at RT with agitation. Blot was incubated in the secondary antibody provided by AgriSera ([AS09 602](#)) diluted to 1:20 000 in for 1h at RT with agitation. The blot was washed as above and developed for 1 min with ECLplus according to the manufacturers instructions. Exposure time was 5 min.

Courtesy of Kaori Takahashi at Hokkaido University, Japan