

This product is **for research use only** (not for diagnostic or therapeutic use)

contact: support@agrisera.com

Agrisera AB | Box 57 | SE-91112 Vännäs | Sweden | +46 (0)935 33 000 | www.agrisera.com

Product no AS16 3140

Anti-ACT 2,8,11 | Actin 2, 8, 11 (clone mAb13a)

Product information

Immunogen	Full-length recombinant <i>Arabidopsis thaliana</i> Actin-2 (ACT2) expressed in <i>E.coli</i> , Uniprot: Q96292
Host	Mouse
Clonality	Monoclonal
Subclass/isotype	IgG1
Purity	IgG1 immunoglobulin purified on Protein G in 0.1 M Sodium Phosphate, pH 7.4, 0.15 M NaCl, 0.05% (w/v) sodium azide.
Format	Liquid
Quantity	100 µg
Storage	Store at -20°C, Make aliquots to avoid repeated freeze-thaw cycles. Please remember to spin the tubes briefly prior to opening them to avoid any losses that might occur from material adhering to the cap or sides of the tube.
Additional information	This antibody is a total IgG fraction, purified on a protein G column. It recognizes actin isoforms ACT2, 8, and 11.

Application information

Recommended dilution	1-2 µg/ml (WB)
Expected apparent MW	45 45 kDa
Confirmed reactivity	<i>Arabidopsis thaliana</i> , <i>Nicotiana tabacum</i> !!AIR8!! <i>Arabidopsis thaliana</i> , <i>Nicotiana tabacum</i>
Predicted reactivity	<i>Actinidia deliciosa</i> , <i>Brachypodium sylvaticum</i> , <i>Brassica napus</i> , <i>Brassica oleracea</i> , <i>Cucumis sativus</i> , <i>Dionaea muscipula</i> , <i>Euphorbia lathyris</i> , <i>Ficus microcarpa</i> , <i>Gentilisea aurea</i> , <i>Gossypium</i> sp., <i>Hevea brasiliensis</i> , <i>Medicago sativa</i> , <i>Medicago truncatula</i> , <i>Oryza sativa</i> subsp. <i>japonica</i> , <i>Oxytropis ochrocephala</i> , <i>Panax notoginseng</i> , <i>Populus trichocarpa</i> , <i>Phaseolus vulgaris</i> , <i>Prunus avium</i> , <i>Striga asiatica</i> , <i>Theobrom cacao</i> , <i>Torenia fournieri</i> , <i>Trifolium pratense</i> , <i>Vaccinium ashei</i> , <i>Ziziphus jujuba</i>
	Species of your interest not listed? Contact us
Not reactive in	No confirmed exceptions from predicted reactivity are currently known
Additional information	Exact working dilution needs to be determined by end user
Selected references	<p>Kandasamy, M.K., et al. (2012). Plant vegetative and animal cytoplasmic actins share functional competence for spatial development with protists. <i>Plant Cell</i> May;24(5):2041-57. doi: 10.1105/tpc.111.095281.</p> <p>Kandasamy, M.K., et al. (2001). One plant actin isoform, ACT7, is induced by auxin and required for normal callus formation. <i>Plant Cell</i>. Jul;13(7):1541-54.</p> <p>Kandasamy, M.K., et al. (1999). The late pollen-specific actins in angiosperms. <i>Plant Journal</i>. Jun;18(6):681-9.</p>