

Product no **AS09 527****Anti-AGO1 | Argonaute 1****Product information****Immunogen** | KLH-conjugated, N-terminal peptide of *Arabidopsis thaliana* AGO1 [O04379](#), [At1q48410](#)**Host** | Rabbit**Clonality** | Polyclonal**Purity** | Immunogen affinity purified serum in PBS pH 7.4.**Format** | Lyophilized**Quantity** | 50 µg**Reconstitution** | For reconstitution add 50 µl of sterile water**Storage** | Store lyophilized/reconstituted at -20°C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please remember to spin the tubes briefly prior to opening them to avoid any losses that might occur from material adhering to the cap or sides of the tube.**Additional information** | Antibody binds microRNA and tasiRNAs, preference for 21nt miRNAs with 5'U.To detect AGO1 in *Nicotiana benthamiana*, please [inquire](#).Recommended for detection of AGO1: [extreme low femtogram range chemiluminescent detection reagent](#)**Application information****Recommended dilution** | 2 µg (ChIP), 1: 200 (IF), (IL), small-RNA-IP-Seq, 1 : 5000-1 : 10 000 (WB)**Expected | apparent MW** | 116.4 | 130 kDa**Confirmed reactivity** | *Arabidopsis thaliana*, *Hyacinthus orientalis*, *Nicotiana benthamiana***Predicted reactivity** | *Brassica pekinensis*, *Capsella rubella*, *Cucumis sativus*, *Glycine max*, *Malus domestica*, *Pisum sativum*, *Ricinus communis*, *Solanum lycopersicum*, *Solanum tuberosum*, *Vitis vinifera*Species of your interest not listed? [Contact us](#)**Not reactive in** | *Chlamydomonas reinhardtii*, *Medicago sativa*, *Salvia sp.*, *Spinacia oleracea*, *Oryza sativa*, *Phaseolus vulgaris*, *Triticum aestivum*, *Zea mays*

Mammals

**Additional information** | AGO expression may be tissue specific and using floral tissue is recommended where most of the AGOs are expressed the highest. Use of proteasome inhibitors as MG132 can help to stabilize AGO proteins during extraction procedure. Buffer for extraction of AGO proteins: [Paudel et al. 2018](#).

The AGO1 antibody is extremely specific to AGO1 and does not cross-react with other antibodies. The evidence is 1) the peptide to which it was raised is at the very N-terminus of the protein and is not present in other AGOs 2) aAGO1 does not cross react with the AGOs which are overexpressed (AGO2, AGO3, AGO4, AGO5, AGO6, AGO9) using a western blot.

[TCA acetone precipitation method](#)**Selected references** | [Bressendorff et al. \(2025\)](#). Importance of an N-terminal structural switch in the distinction between small RNA-bound and free ARGONAUTE. *Nat Struct Mol Biol.* 2025 Jan 7. doi: 10.1038/s41594-024-01446-9.  
[Nasfi et al. \(2024\)](#). A pipeline for validation of *Serendipita indica* effector-like sRNA suggests cross-kingdom communication in the symbiosis with *Arabidopsis*. *J Exp Bot.* 2024 Dec 26;erae515. doi: 10.1093/jxb/erae515.  
[Li et al. \(2024\)](#). RNA helicase Brr2a promotes miRNA biogenesis by properly remodelling secondary structure of pri-miRNAs. *Nat Plants.* 2024 Oct;10(10):1532-1547. doi: 10.1038/s41477-024-01788-8.  
[Mirlohi et al. \(2024\)](#). An AGO10: miR165/6 module regulates meristem activity and xylem development in the *Arabidopsis* root. *EMBO J.* 2024 May;43(9):1843-1869. doi: 10.1038/s44318-024-00071-y.  
[Martín-Merchán et al. \(2024\)](#). *Arabidopsis* AGO1 N-terminal extension acts as an essential hub for PRMT5 interaction and post-translational modifications. *Nucleic Acids Res.* 2024 May 20;gkae387. doi: 10.1093/nar/gkae387.  
[Barre-Villeneuve et al. \(2024\)](#). The unique dual targeting of AGO1 by two types of PRMT enzymes promotes phasiRNA loading in *Arabidopsis thaliana*. *Nucleic Acids Res.* 2024 Mar 21;52(5):2480-2497. doi: 10.1093/nar/gkae045.  
[Blagojevic et al. \(2024\)](#). Heat stress promotes *Arabidopsis* AGO1 phase separation and association with stress granule components. *iScience.* 2024 Feb 6;27(3):109151. doi: 10.1016/j.isci.2024.109151. eCollection 2024 Mar 15.  
[Bradamante et al. \(2023\)](#). Two ARG ONAUTE proteins loaded with transposon-derived small RNAs are associated with

the reproductive cell lineage in Arabidopsis. Plant Cell. 2023 Dec 7;:koad295.doi: 10.1093/plcell/koad295.


Meng et al. (2022) The novel activity of Argonautes in intron splicing: A transcriptome-wide survey in plants. J Plant Physiol. 2022 Jan 31;270:153632. doi: 10.1016/j.jplph.2022.153632. Epub ahead of print. PMID: 35114616.

Cabezas-Fuster et al. (2022). Missplicing suppressor alleles of Arabidopsis PRE-MRNA PROCESSING FACTOR 8 increase splicing fidelity by reducing the use of novel splice sites. Nucleic Acids Res. 2022 Jun 10;50(10):5513-5527. doi: 10.1093/nar/gkac338. PMID: 35639749; PMCID: PMC9177961.

Delenko et al. (2022) MicroRNA biogenesis and activity in plant cell dedifferentiation stimulated by cell wall removal. BMC Plant Biol. 2022 Jan 3;22(1):9. doi: 10.1186/s12870-021-03323-9. PMID: 34979922; PMCID: PMC8722089. (immunofluorescence)

Oliver & Martinez. (2021) Accumulation dynamics of ARGONAUTE proteins during meiosis in Arabidopsis. Plant Reprod. 2021 Nov 23. doi: 10.1007/s00497-021-00434-z. Epub ahead of print. PMID: 34812935.

Dalmadi et al. (2021) Controlled RISC loading efficiency of miR168 defined by miRNA duplex structure adjusts ARGONAUTE1 homeostasis. Nucleic Acids Res. 2021 Dec 16;49(22):12912-12928. doi: 10.1093/nar/gkab1138. PMID: 34850097; PMCID: PMC8682782.



80 µg of *Arabidopsis thaliana* soluble total cell extract (extracted in 20 mM Tris pH 7.5, 5mM MgCl<sub>2</sub>, 2.5mM DTT, 300 mM NaCl, 0.1% NP-40, 1% proteasome inhibitor MG132) was separated on 6% SDS-PAGE and blotted 1h to nitrocellulose. Filters were blocked 1h with 5% low-fat milk powder in TBS-TT (0.25% TWEEN20; 0.1% Triton-X) and probed with anti-AGO1 antibody (1:10 000, 1h) and secondary anti-rabbit (1:10000, 1 h) antibody (HRP conjugated) in TBS-TT containing 5% low fat milk powder. Antibody incubations were followed by washings in TBS-TT. All steps were performed at RT with agitation. Blots were developed for 5 min with ECL-PLUS detection reagent according to the manufacturer's instructions. Exposure time was 5 seconds.

Roche's protease inhibitor cocktail (no EDTA) can also be applied in extraction buffer.

Courtesy Dr. Ericka Havecker, University of Cambridge, UK