


Product no **AS14 2797****Anti-AGO1-PAZ | Argonaute 1 PAZ domain (Chlamydomonas)****Product information**

| | |
|-----------------------|---|
| Immunogen | KLH-conjugated peptide derived from AGO1-PAZ domain of <i>Chlamydomonas reinhardtii</i> Cre02.g141050.t1.1 |
| Host | Rabbit |
| Clonality | Polyclonal |
| Purity | Immunogen affinity purified serum in PBS pH 7.4. |
| Format | Lyophilized |
| Quantity | 50 µg |
| Reconstitution | for reconstitution add 50 µl of sterile water |
| Storage | Store lyophilized/reconstituted at -20°C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please remember to spin the tubes briefly prior to opening them to avoid any losses that might occur from material adhering to the cap or sides of the tube. |

Application information

| | |
|-------------------------------|---|
| Recommended dilution | 1: 500 (WB) |
| Expected apparent MW | 109 120 kDa |
| Confirmed reactivity | <i>Chlamydomonas reinhardtii</i> |
| Additional information | AGO1 signal is detected at higher MW most likely due to post-translational modifications) |

Application example

Total proteins of *Chlamydomonas reinhardtii* protein saturated in 8M urea were separated on 15% SDS-PAGE and blotted for 1 hour to 0.2 µm nitrocellulose at 100V using wet transfer system. Blots were blocked with 0.5% cold fish gelatin for 1hr at room temp with agitation. Blot was incubated in the primary antibody (anti-H3) at a dilution of 1:2500 for an hour at RT with agitation. The blots were washed with 3X 15min TBS-TT at RT with agitation. Blots as incubated in the secondary antibody (goat anti-rabbit IgG (H&L), DyLight® 800 conjugated) 1:2500 dilution for 30min at RT with agitation and washed 1X with TBSTT for 15min, 1X with TBST for 15min before scanning with the ODyssey IRD scanner.

Courtesy of Dr. Betty Chung, University of Cambridge, United Kingdom