

This product is for research use only (not for diagnostic or therapeutic use)

contact: support@agrisera.com

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | www.agrisera.com

Product no AS22 4789

AHA2 p | ATPase 2 phosphorylated, plasma membrane-type

Product information

Immunogen KLH-conjugated peptide derived from Arabidopsis thaliana AHA2 protein sequence UniProt: P19456, TAIR: At4g30190

Host Rabbit

Clonality Polyclonal

Purity Antigen affinity purified serum, in PBS pH 7.4

Format Lyophilized

Quantity 50 μg

Reconstitution For reconstitution, add 50 μl, of sterile or deionized water.

Storage Store lyophilized/reconstituted at -20 °C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please, remember to spin tubes briefly prior to opening them to avoid any losses that might occur from lyophilized

material adhering to the cap or sides of the tubes.

Application information

Recommended dilution 1:10 000 (WB)

Expected | apparent


104,4 | kDa

Predicted reactivity Brassica oleracea, Camelina sativa, Capsella rubella, Eutrema salsugineum, Raphanus sativus Species of your interest

not listed? Contact us

Not reactive in No confirmed exceptions from predicted reactivity are currently known

Selected references To be added when available, antibody available in October 2023.



Samples:

- 1 Recombinant protein MBP-C terminal region of AHA2, phosphorylation treatment
- 2 Recombinant protein MBP-C terminal region of AHA2, subsequently dephosphorylated
- 3 Recombinant protein MBP, phosphorylation treatment
- 4 Recombinant protein MBP-C terminal region of AHA2, Thr947Ala, phosphorylation treatment 5- Mark: MW markers

1 μg/well of recombinant protein produced in *E. coli* and purified by affinity chromatography was incubated in the reaction mixture (30 ul). 3 ul of the reaction were mixed with 6 ul Laemli 2X and denatured at 95°C for 5 min. Samples were separated on 10% SDS-PAGE and blotted overnight (ON) to PVDF (Inmobilon®-FL) (pore size of 0.45 μm) using wet transfer. The blot was blocked with 3 % non-fat milk 2h/RT with agitation. The blot was incubated in the primary antibody at a dilution of 1:10 000 for 90 min/RT with agitation 3% non-fat milk TBS with agitation. The antibody solution was decanted, and the blot was rinsed briefly twice, then washed once for 15 min and 3 times for 5 min in TBS-T at RT with agitation. The blot was incubated in a matching secondary antibody (anti-rabbit IgG horseradish peroxidase conjugated) diluted to 1: 5000 for 90 min/RT with agitation. The blot was washed as above and developed with the following chemiluminescent detection reagent: AgriseraBright. Exposure time was 2 min.


Courtesy of Cristian Mayordomo, IBMCP, Universidad Politécnica de Valencia, Spain



This product is for research use only (not for diagnostic or therapeutic use)

contact: support@agrisera.com

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | www.agrisera.com



Titration of the AS22 4789: AHA2 p | ATPase 2 phosphorylated, plasma membrane-type antibody. Ten μ g of protein from Amaranth stem mixed membranes (TO), or from beetroot plasma membranes either pre-phosphorylated, i.e. treated with ATP (MP+P), or not (MP* and MP-P) were blotted and probed with the indicated dilutions of antibody. Molecular mass markers (MPM) have, from top to bottom 250, 130, 100, 70 (red) and 55 kDa, respectively.

Courtesy Dr. Luis González, CINVESTAV, Mexico