


Product no **AS18 4230****Anti-AHB1 | Non-symbiotic hemoglobin 1 (A. thaliana)****Product information**

Immunogen	KLH-conjugated peptide derived from <i>Arabidopsis thaliana</i> AHB1, UniProt: O24520 , TAIR: AT2G16060
Host	Rabbit
Clonality	Polyclonal
Purity	Antigen affinity purified serum, in PBS pH 7.4.
Format	Lyophilized
Quantity	50 µg
Reconstitution	For reconstitution add 50 µl, of sterile water.
Storage	Lyophilized antibody can be stored at -20°C for up to 3 years. Once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please, remember to spin tubes briefly prior to opening them to avoid any losses that might occur from lyophilized material adhering to the cap or sides of the tubes.

Application information

Recommended dilution	1 : 1000 (WB)
Expected apparent MW	kDa
Confirmed reactivity	<i>Arabidopsis thaliana</i>
Predicted reactivity	<i>Malus domestica</i>
Not reactive in	No confirmed exceptions from predicted reactivity are currently known
Additional information	Conditions for detection of endogenous AHB1 remain to be determined.
Selected references	To be added when available. Antibody released in April 2022.

**Samples:**

- 1 - 90 µg of *Arabidopsis thaliana* whole seedling extract, Col-0 (wild type) submerged to induce PGB1
- 2 - 90 µg of *Arabidopsis thaliana* whole seedling extract, Col-3 (wild type) submerged to induce PGB1
- 3 - 90 µg of *Arabidopsis thaliana* whole seedling extract, Col-0; 35S: GFP-PGB1 in air (control)
- 4 - 90 µg of *Arabidopsis thaliana* whole seedling extract, C24; 35S:PGB1 in air (control)
- 5 - 90 µg of *Arabidopsis thaliana* whole seedling extract, C24; 35S:PGB1 submerged to induce PGB1
- 6 - 90 µg of *Arabidopsis thaliana* whole seedling extract, Col-0 (wild type in air (control)
- 7 - 90 µg of *Arabidopsis thaliana* whole seedling extract, C24 (wild type) in air (control)
- 8 - 90 µg of *Arabidopsis thaliana* whole seedling extract, knock down mutant of PGB1 (C24 background) submerged to induce PGB1
- 9 - 90 µg of *Arabidopsis thaliana* whole seedling extract, knock out mutant of PGB1 (Col-3 background) submerged to induce PGB1 (AHB1)

Mark: MW markers: Dual Color (BioRad)

90 µg/well of total protein extracted freshly from *Arabidopsis thaliana* seedlings with extraction buffer (100mM Tris-HCl, 150mM NaCl, 0.25% NP-40) containing 1mM PMSF and 1X cOmplete® EDTA-free proteases inhibitors, Sigma) and denatured with buffer (Tris-HCl 0,25 M, pH 6,8; SDS 8% (w/v); glycerol 40% (v/v); -Mercaptoethanol 20% (v/v) and Bromophenol Blue) at 90°C for 10 min were separated on 12% SDS-PAGE and blotted (20min; 2A; 25V) to PVDF (pore size of 0.2 µm), using semi-dry transfer. Blot was blocked with 5% milk for 1h/RT with agitation. Blot was incubated in the primary antibody at a dilution of 1: 1000 ON/4°C with agitation. The antibody solution was decanted and the blot was washed 2 times for 5 min in TBS-T at RT with agitation. Blot was incubated in Agrisera matching secondary antibody (Goat anti-rabbit IgG HRP peroxidase conjugated, [AS09 602](#)) diluted to 1:25 000 in for 1h/RT with agitation. The blot was washed 4 times for 5 min in TBS-T at RT with agitation and developed with Agrisera ECL SuperBright, [AS16 ECL-S](#). Exposure time was 5 seconds.

The lines used in the experiments were generated by:

This product is **for research use only** (not for diagnostic or therapeutic use)

contact: support@agrisera.com

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | www.agrisera.com

-[Hebeslstrup et al., 2008](#) corresponds to 35S: GFP-PGB1
-[Perazzoli et al., 2004](#) corresponds to 35S:PGB1

Courtesy of Dr. Inmaculada Sánchez-Vicente and Isabel Manrique-Gil, USAL-CIALE Salamanca, Spain