

This product is for research use only (not for diagnostic or therapeutic use)

contact: support@agrisera.com

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | www.agrisera.com

Product no AS09 462

Anti-AKINB2 | SNF1-related protein kinase regulatory subunit beta-2

Product information

Immunogen KLH-conjugated peptide derived from Arabidopsis thaliana AKIN beta-2 Q9SCY5

Host Rabbit

Clonality Polyclonal

Purity Immunogen affinity purified serum in PBS pH 7.4.

Format Lyophilized

Quantity 200 μg

Reconstitution For reconstitution add 200 μl of sterile water

Storage Storage Store lyophilized/reconstituted at -20 °C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please remember to spin the tubes briefly prior to opening them to avoid any losses that might occur from material adhering to

the cap or sides of the tube.

Application information

Recommended dilution 2 μg/ml

Expected | apparent


31.9 | 35 kDa

Not reactive in No confirmed exceptions from predicted reactivity are currently known

Selected references Emanuelle et al. (2015). SnRK1 from Arabidopsis thaliana is an atypical AMPK. Plant J. 2015 Mar 3. doi:

10.1111/tpj.12813.

Application example



25 μg of empty vector (1), pCDNA 3.1 – HA- tagged AKIN-beta1 (2), pCDNA 3.1 – HA- tagged AKIN-beta2 (3), 50 μg of total protein from *Arabidopsis thaliana* leaves (4) had been homogenized into 50 mM NaPO4 pH 7.5, 20 mM KCl, 0.5 M sucrose, 0.2 mM PMSF, 10 mM DTT and protease inhibitor cocktail (Sigma, P9599). Extracts were separared on 10%NuPage Bis-Tris Novex SDS PAGE (Invitrogen) gels followed by a transfer for 60 min to PVDF membrane. Filters were blocked for 1h with 5% low-fat milk powder in PBS-T (0.1% TWEEN 20) and probed with anti-AKINbeta-2 antibodies (2 μg/ml for 60 min at RT) followed by Protein G-HRP (1:3 000 for 60 min at RT) in PBS-T. Antibody incubations were followed by washings in PBS-T (3 x 5 min). All washing steps were performed at RT with agitation. Signal was detected with chemiluminescence, using CCD camera. Exposure time was 10 seconds.

Courtesy Dr. David Stapleton, Melbourne, Australia