


Product no **AS23 4921****Anti-NAB1 | nucleic acid binding protein 1, Chlamydomonas****Product information****Immunogen** | KLH-conjugated peptide derived from *Chlamydomonas reinhardtii* NAB1 protein sequence, UniProt [Q8GV23](#)**Host** | Rabbit**Clonality** | Polyclonal**Purity** | Antigen affinity purified serum, in PBS pH 7.4**Format** | Lyophilized**Quantity** | 50 µg**Reconstitution** | For reconstitution, add 50 µl of sterile or deionized water.**Storage** | Store lyophilized/reconstituted at -20°C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please, remember to spin tubes briefly prior to opening them to avoid any losses that might occur from lyophilized material adhering to the cap or sides of the tubes.**Additional information** | The NAB1 gene ID in Phytozome is Cre06.g268600_45321. There are two transcripts: Cre06.g268600.4532.1 - - 35,324 kDa and Cre06.g268600.4532.2 - - 26,539 kDa. In cell bodies of gametes, produce proteins from both transcripts, while in vegetative cells only one transcript is translated to NAB1 protein (26.5 kDa).**Application information****Recommended dilution** | 1 : 1000 (WB)**Expected | apparent MW** | 26.5 kDa | 26 kDa (for *Chlamydomonas reinhardtii*)**Confirmed reactivity** | *Chlamydomonas reinhardtii***Predicted reactivity** | Species of your interest not listed? [Contact us](#)**Not reactive in** | No confirmed exceptions from predicted reactivity are currently known**Selected references** | To be added when available, antibody available in October 2024.**Samples:**1 - 2 ug of *C. reinhardtii* C1 whole extract2 - 2 ug of *C. reinhardtii* C1 cell body3 - 2 ug of *C. reinhardtii* C1 cilia Mark: MW markers 4 - 2 ug of *C. reinhardtii*

Chlamydomonas cilia isolation was performed, and whole cell extract, cell body, and cilia fractions were stored at -80°C. The buffer components consisted of 20 mM HEPES, 10 mM MgCl₂, 50 mM KCl, 2 mM DTT, 1 mM EDTA, and 1X Protease inhibitor. Samples were denatured with 6x SDS-Sample Laemmli buffer at 95°C for 5 minutes and store at -20 °C. Samples were separated on 10% SDS-PAGE and transferred overnight to PVDF membranes (pore size of 0.45 µm) using wet transfer in cold conditions. The blot was then blocked with 5% milk for 1 hour at room temperature with agitation. Next, the blot was incubated in the primary antibody at a dilution of 1:1000 for 1 hour at room temperature in TBS-T with agitation. Subsequently, the antibody solution was decanted, and the blot was briefly rinsed twice, followed by three washes for 10 minutes each in TBS-T at room temperature with agitation. The blot was then incubated with goat anti-rabbit IgG secondary antibody (Agrisera goat anti-rabbit IgG horse radish peroxidase conjugated [AS09 602](#)) at a dilution of 1:25000 for 1 hour at room temperature with agitation. After

incubation with the secondary antibody, the blot was washed three times as previously described and developed using a chemiluminescent detection reagent with exposure time of 30 seconds.

Courtesy of Meenakshi Tanwar, Department of Cell Biology and Molecular Genetics, University of Maryland, USA