


Product no **AS23 4929****Anti-TBP1/2 | TATA BINDING PROTEIN 1/2****Product information**

Immunogen	KLH-conjugated peptide derived from protein sequence of <i>Arabidopsis thaliana</i> TBP1 and TBP2 UniProt: P28147 and P28148 , TAIR: AT3G13445.1 , AT1G55520.1
Host	Rabbit
Clonality	Polyclonal
Purity	Antigen affinity purified serum, in PBS pH 7.4
Format	Lyophilized
Quantity	50 µg
Reconstitution	For reconstitution, add 50 µl, of sterile or deionized water.
Storage	Store lyophilized/reconstituted at -20 °C; once reconstituted, make aliquots to avoid repeated freeze-thaw cycles. Please, remember to spin tubes briefly prior to opening them to avoid any losses that might occur from lyophilized material adhering to the cap or sides of the tubes.

Application information

Recommended dilution	6 µg
Expected apparent MW	22.4 kDa
Confirmed reactivity	<i>Arabidopsis thaliana</i>
Predicted reactivity	<i>Arachis hypogaea</i> , <i>Brachypodium distachyon</i> , <i>Brassica napus</i> , <i>Cannabis sativa</i> , <i>Capsicum annuum</i> , <i>Glycine max</i> , <i>Hordeum vulgare</i> , <i>Malus domestica</i> , <i>Manihot esculenta</i> , <i>Medicago truncatula</i> , <i>Nicotiana tabacum</i> , <i>Oryza sativa</i> , <i>Pisum sativum</i> , <i>Solanum lycopersicum</i> , <i>Solanum tuberosum</i> , <i>Sorghum bicolor</i> , <i>Spinacia oleracea</i> , <i>Theobroma cacao</i> L, <i>Triticum sp.</i> , <i>Vitis vinifera</i> , <i>Zea mays</i>
	Species of your interest not listed? Contact us
Not reactive in	No confirmed exceptions from predicted reactivity are currently known
Additional information	The peptide used to elicit this antibody is conserved in both isoforms: 1 and 2 of TATA Binding Protein.
Selected references	To be added when available, antibody available released in October 2025.



Chromatin extraction and immunoprecipitation was performed according to the aChIP protocol (<https://doi.org/10.1038/s41477-024-01743-7>). In particular, all the buffer formulations and wash schemes were preserved. In brief, ground in liquid nitrogen powder from 0.2g of green siliques of *Arabidopsis thaliana*, was crosslinked for 10 min in 4 °C in 20ml of MC buffer (0.1M sucrose, 10mM sodium phosphate pH 7.0, 50mM NaCl) containing 1% formaldehyde. After stopping the crosslink with 0.125 M glycine, the pellet was resuspended in Z-1 buffer. Single washes using Z-2 and Z-3 were performed and the chromatin was resuspended in 2 ml of Z-4. Samples were sonicated using Bioruptor (Diagenode) apparatus for 26 cycles (30sec ON/30sec OFF; Mode: High). The average fragment size was around 500bp, estimated from agarose gel migration. After sonication and 10 min centrifugation, the soluble supernatant was collected and directly used for IP. Before, 12 µl of Protein-G-Dynabeads (Thermo Sci) were incubated in PBS supplemented with 0.5% BSA and with or without 6µg of anti-TBP antibodies (Agrisera) for 4 hrs in 4 °C with constant rotation, prior to adding to chromatin. Immunoprecipitation reactions were set for overnight. Each chromatin sample was split to 2 IP reactions (TBP and NoAb, 600µl each) and 1 Input sample (60µl). On the next day, IP reactions were washed twice using Low Salt, once using



This product is **for research use only** (not for diagnostic or therapeutic use)

contact: support@agrisera.com

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | www.agrisera.com

High Salt followed by LiCl Wash solutions, then twice using TE buffer. DNA was recovered from IP and Input samples by 15 min incubation in 10% chelex suspension in 95°C, followed by proteinase K digestion (2h in 50°C with 2µl of Proteinase K) and inactivation (10 min in 95°C). qPCR was performed on IP samples, No Antibody controls and 1:10 diluted Input samples (1:100 - total dilution factor, relative to IP), using SYBR I qPCR mix (Roche).

Courtesy of Dr. Lien Brzeźniak, Polish Academy of Science, Warsaw, Poland