

This product is for research use only (not for diagnostic or therapeutic use)

contact: support@agrisera.com

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | www.agrisera.com

## Product no AS14 2816

## Anti-ASH1 | Histone-lysine N-methyltransferase ASH1

## **Product information**

Immunogen N-terminal GST-fusion of the peptide containing amino acids 1756-1855 of the Drosophila melanogaster Ash1protein,

UniProt: Q9VW15

**Host** Rabbit

Clonality Polyclonal

**Purity** Immunogen affinity purified serum in PBS pH 7.4.

Format Lyophilized

Quantity 50 μg

**Reconstitution** For reconstitution add 50 μl of sterile water

Storage Store lyophilized/reconstituted at -20°C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please remember to spin the tubes briefly prior to opening them to avoid any losses that might occur from material adhering to

the cap or sides of the tube.

## **Application information**

Recommended dilution 1:500 (IL), 10ng/ul (ChIP)

Expected | apparent 236 | 300 kDa

MW

Confirmed reactivity Drosophila melanogaster

Predicted reactivity Drosophila melanogaster

Not reactive in No confirmed exceptions from predicted reactivity are currently known

Selected references Kahn et al. (2016). Interdependence of PRC1 and PRC2 for recruitment to Polycomb Response Elements. Nucleic

Acids Res. 2016 Aug 23. pii: gkw701. [Epub ahead of print].

Lee et al. (2015). Genome-wide activities of Polycomb complexes control pervasive transcription. Genome Res. 2015 Aug;25(8):1170-81. doi: 10.1101/gr.188920.114. Epub 2015 May 18.