

This product is for research use only (not for diagnostic or therapeutic use)

contact: support@agrisera.com

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | www.agrisera.com

Product no AS22 4718

AURKAIP1 mouse | Aurora kinase A-interacting protein

Product information

Immunogen Recombinat mouse AURKAIP1 protein expressed in E.coli, UniProt: Q9DCJ7

Host Rabbit

Clonality Polyclonal

Purity Antigen affinity purified serum, in PBS pH 7.4

Format Lyophilized

Quantity 50 μg

Reconstitution For reconstitution, add 50 μl, of sterile or deionized water.

Storage Store lyophilized/reconstituted at -20 °C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please, remember to spin tubes briefly prior to opening them to avoid any losses that might occur from lyophilized

material adhering to the cap or sides of the tubes.

Application information

Recommended dilution 1:500 (WB)

Expected | apparent

22 | 17 kDa (without mitochondrial target sequence MTS)


Confirmed reactivity Mouse

Predicted reactivity | Species of your interest not listed? Contact us

Not reactive in No confirmed exceptions from predicted reactivity are currently known

Additional information This antibody is also recognizing human AURKAIP1.

Selected references To be added when available, antibody available in November 2022.



10 µg of mouse mitochondrial lysate extracted with RIPA buffer (150 mM sodium chloride, 1.0% NP-40, 0.5% sodium deoxycholate 0.1% SDS, 50 mM Tris, pH 8.0) and denatured in NuPAGE LDS Sample Buffer (Invitrogen, NP0007) at 70°C for 5 min were separated in 4-12% NuPAGE Bis-Tris (Invitrogen, NP0329BOX) and blotted 7min to Nitrocellulose using the iBlot transfer system (Invitrogen). Blots were blocked with 5% milk at RT 30 min. Primary antibodies were used at 1: 500 ON/4°C incubaftion, following with the 3x15 min,. wash with TBS-T at RT. Secondary antibodies were diluted to 1: 25 000, incubated for 1h/RT (AS09 602, Agrisera). Reaction was developed with Agrisera ECLSuperBright (AS16 ECL-S-10).

Courtesy of Dr.

Rodolfo Garcia Villegas, Karolinksa Institute, Sweden