

This product is **for research use only** (not for diagnostic or therapeutic use)

contact: support@agrisera.com

Agrisera AB | Box 57 | SE-911121 Vännäs | Sweden | +46 (0)935 33 000 | www.agrisera.com

Product no AS09 481-DL650

Anti-BiP | Lumenal-binding protein, DyLight® 650 conjugated (40 µg)

Product information

Immunogen	KLH-conjugated synthetic peptide derived from <i>Arabidopsis thaliana</i> BiP proteins: BiP1 At5g28540 Q9LKR3 , BiP2 At5g42020 F4K007 , BiP3 At1g09080 Q8H1B3
Host	Rabbit
Clonality	Polyclonal
Purity	Immunogen affinity purified serum, in PBS pH 7.4, conjugated to DyLight® 650.
Format	Liquid in PBS pH 7.4.
Quantity	40 µg
Storage	Store at 4°C for 12-18 months, A preservative may be added for long time storage up to 2 years. Spin briefly before use.
Additional information	DyLight® 650 has Amax = 652 nm, Emax = 672 nm. DyLight® is a registered trademark of Thermo Fisher Inc., and its subsidiaries.

Application information

Recommended dilution	To be determined by end user
Expected apparent MW	73.5 80 kDa
Confirmed reactivity	<i>Arabidopsis thaliana</i> , <i>Brassica napus</i> , <i>Chlamydomonas reinhardtii</i> , <i>Cucumis sativus</i> , <i>Nicotiana benthamiana</i> , <i>Raphanus sativa L</i> , <i>Tokinashi-daiikon</i> , <i>Olea europaea</i> , <i>Picea abies</i> , <i>Physcomitrella patens</i> , <i>Spinacia oleracea</i> , <i>Solanum lycopersicum</i> , <i>Solanum tuberosum</i> , <i>Triticum aestivum</i> , <i>Zea mays</i>
Predicted reactivity	<i>Arabis alpina</i> , <i>Capsella rubella</i> , <i>Capsicum annuum</i> , <i>Citrus clementina</i> , <i>Citrus sinsensis</i> , <i>Eucalyptus grandis</i> , <i>Glycine max</i> , <i>Hordeum vulgare</i> , <i>Isatis tinctoria</i> , <i>Prunus persica</i> , <i>Triticum aestivum</i> , <i>Petunia hybrida</i> , <i>Picea sitchensis</i> , <i>Populus trichocarpa</i> , <i>Ricinus communis</i> , <i>Vitis vinifera</i>
	Species of your interest not listed? Contact us
Not reactive in	No confirmed exceptions from predicted reactivity are currently known.
Additional information	Protein or membrane sample should be treated at 70°C for 10 min before loading on the gel, This antibody has so far not worked in IP.
Selected references	To be added when available. Antibody released in May 2023.