


Product no **AS14 2760****Anti-CsoS1A/B/C | Major carboxysome shell protein 1A, AB, 1C****Product information**

|                       |   |
|-----------------------|---|
| <b>Immunogen</b>      | KLH-conjugated peptide conserved in Major carboxysome shell protein 1A, UniProt: <a href="#">P45689</a> , Major carboxysome shell protein 1B UniProt: <a href="#">P45690</a> , Major carboxysome shell protein 1C, UniProt: <a href="#">P45688</a>                            |
| <b>Host</b>           | Rabbit  |
| <b>Clonality</b>      | Polyclonal  |
| <b>Purity</b>         | Immunogen affinity purified serum in PBS pH 7.4.  |
| <b>Format</b>         | Lyophilized   |
| <b>Quantity</b>       | 50 µg   |
| <b>Reconstitution</b> | For reconstitution add 50 µl of sterile water   |
| <b>Storage</b>        | Store lyophilized/reconstituted at -20°C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please remember to spin the tubes briefly prior to opening them to avoid any losses that might occur from material adhering to the cap or sides of the tube. |

**Application information**

|                               |   |
|-------------------------------|---|
| <b>Recommended dilution</b>   | 1 : 5000 (WB)   |
| <b>Expected   apparent MW</b> | 9-11 kDa  |
| <b>Confirmed reactivity</b>   | <i>Halothiobacillus neapolitanus</i> (strain ATCC 23641 / c2) ( <i>Thiobacillus neapolitanus</i> )  |
| <b>Predicted reactivity</b>   | Algae (red), Cyanobacteria, Cryptomonads<br>Species of your interest not listed? <a href="#">Contact us</a>   |
| <b>Not reactive in</b>        | <i>Synechococcus elongatus</i>  |
| <b>Selected references</b>    | <p><a href="#">Li et al. (2024)</a>. Nanoengineering Carboxysome Shells for Protein Cages with Programmable Cargo Targeting. ACS Nano. 2024 Mar 12;18(10):7473-7484. doi: 10.1021/acsnano.3c11559.</p> <p><a href="#">Huang, Jiang, Yang, et al. (2022)</a> Probing the Internal pH and Permeability of a Carboxysome Shell. Biomacromolecules. 2022;23(10):4339-4348. doi:10.1021/acs.biomac.2c00781</p> <p><a href="#">Chen et al. (2022)</a> ACS Synth. Biol. 2022, 11, 1, 154-161 Publication Date: October 19, 2021 <a href="https://doi.org/10.1021/acssynbio.1c00311">https://doi.org/10.1021/acssynbio.1c00311</a></p> <p><a href="#">Li et al. (2020)</a>. Reprogramming bacterial protein organelles as a nanoreactor for hydrogen production. Nat Commun. 2020 Oct 28;11(1):5448. doi: 10.1038/s41467-020-19280-0. PMID: 33116131; PMCID: PMC7595155.</p> <p><a href="#">Long et al. (2018)</a>. Carboxysome encapsulation of the CO<sub>2</sub>-fixing enzyme Rubisco in tobacco chloroplasts. Nat Commun. 2018 Sep 3;9(1):3570. doi: 10.1038/s41467-018-06044-0.</p> |



0.125 - 1.0 µg of total purified *Halothiobacillus neapolitanus* carboxysome protein (equivalent to approximately 20 - 170 ng CsoS1A, B and C proteins) were separated on 4-20 % Bio-Rad TGX stain-free SDS-PAGE and blotted 1h to PVDF using a Bio-Rad TransBlot Turbo Blotting System. Blots were blocked with 5% skim milk powder in TBS-T for 1h at room temperature (RT) with agitation. Blot was incubated in the primary antibody at a dilution of 1:5 000 for 1h at RT with agitation. The antibody solution was decanted and the blot was rinsed 3 times for 5 min in TBS-T at RT with agitation. Blot was incubated in secondary antibody (goat-anti-rabbit IgG alkaline phosphatase conjugated) diluted to 1:5 000 in TBS-T for 1h at RT with agitation. The blot was washed as above and antibody binding determined using the Promega Attophos reagent system as described by the manufacturer. Reaction time was 10 seconds and membranes were imaged using a Bio-Rad VersaDoc.

Courtesy of Dr. Ben Long, The Australian National University, Australia