

Product no **AS18 4169****Anti-Cyt b6 / PetB | Thylakoid membrane cytochrome b6 protein, N terminal****Product information**

Immunogen	KLH-conjugated peptide chosen from <i>Arabidopsis thaliana</i> PetB protein sequence, UniProt: P56773 , TAIR: ATCG00720
Host	Rabbit
Clonality	Polyclonal
Purity	Serum
Format	Lyophilized
Quantity	50 µl
Reconstitution	For reconstitution add 50 µl, of sterile water
Storage	Lyophilized antibody can be stored at -20 °C for up to 3 years. Re-constituted antibody can be stored at 4 °C for several days to weeks. Once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please remember to spin the tubes briefly prior to opening them to avoid any losses that might occur from material adhering to the cap or sides of the tube.

Application information

Recommended dilution	1 : 1000 - 1 : 5000 (WB)
Expected apparent MW	24 kDa
Confirmed reactivity	<i>Arabidopsis thaliana</i> , <i>Chlamydomonas reinhardtii</i> , <i>Echinochloa crus-galli</i> , <i>Marchantia polymorpha</i> , <i>Medicago sativa</i> , <i>Nannochloropsis oceanica</i> strain IMET1, <i>Nicotiana benthamiana</i> , <i>Panicum miliaceum</i> , <i>Pisum sativum</i> , <i>Zea mays</i>
Predicted reactivity	<i>Acacia prairi</i> , <i>Angelphytum bahiense</i> , <i>Aristolochia promissa</i> , <i>Daucus glochidiatus</i> , <i>Dimerostemma brasilianum</i> , <i>Elaphandra paucipunctata</i> , <i>Fraxinus chiisanensis</i> , <i>Gossypium populifolium</i> , <i>Hydrangea heteromalla</i> , <i>Nicotiana sylvestris</i> , <i>Oryza sativa</i> , <i>Salix tetrasperma</i> , <i>Solanum lycopersicum</i> , <i>Thottea siliquosa</i> , <i>Triticum turgidum</i> , <i>Ulva flexuosa</i> , <i>Wedelia biflora</i> , <i>Vanilla aphylla</i> Species of your interest is not listed? Species of your interest not listed? Contact us .
	Species of your interest not listed? Contact us
Selected references	Collombat et al. (2025) . Arabidopsis conditional photosynthesis mutants abc1k1 and var2 accumulate partially processed thylakoid preproteins and are defective in chloroplast biogenesis. Commun Biol . 2025 Jan 22;8(1):111. doi: 10.1038/s42003-025-07497-y. Hani and Krieger-Liszkay (2024) . Manganese deficiency alters photosynthetic electron transport in Marchantia polymorpha. Elsevier Plant Physiology and Biochemistry Available online 16 August 2024, 109042. Sulli et al. (2023) . Generation and physiological characterization of genome edited Nicotiana benthamiana plants containing zeaxanthin as the only leaf xanthophyll. Planta . 2023 Oct 5;258(5):93. doi: 10.1007/s00425-023-04248-3. Zhang et al. (2023) . Bupleurum chinense and Medicago sativa sustain their growth in agrophotovoltaic systems by regulating photosynthetic mechanisms. Renewable and Sustainable Energy Reviews Volume 189, Part B, January 2024, 114024. Nagy et al. (2023) . Photoautotrophic and sustained H2 production by the pgr5 mutant of Chlamydomonas reinhardtii in simulated daily light conditions. International Journal of Hydrogen Energy Volume 53, 31 January 2024, Pages 760-769. Urban, Rogowski & Romanowska (2022) , Crucial role of the PTOX and CET pathways in optimizing ATP synthesis in mesophyll chloroplasts of C3 and C4 plants, Environmental and Experimental Botany, Volume 202, October 2022, 105024, https://doi.org/10.1016/j.envexpbot.2022.105027 Wada et al. (2021) Identification of a Novel Mutation Exacerbated the PSI Photoinhibition in pgr5/pgr1 Mutants; Caution for Overestimation of the Phenotypes in Arabidopsis pgr5-1 Mutant. Cells. 2021 Oct 26;10(11):2884. doi: 10.3390/cells10112884. PMID: 34831107; PMCID: PMC8616342. Chen et al. (2021) Degradation of the photosystem II core complex is independent of chlorophyll degradation mediated by Stay-Green Mg2+ dechelataase in Arabidopsis, Plant Science, Volume 307, 2021, 110902, ISSN 0168-9452, https://doi.org/10.1016/j.plantsci.2021.110902 . Lu et al. (2021) . Role of an ancient light-harvesting protein of PSI in light absorption and photoprotection. Nat Commun. 2021 Jan 29;12(1):679. doi: 10.1038/s41467-021-20967-1. PMID: 33514722; PMCID: PMC7846763. Wang et al. (2020) . Post-translational coordination of chlorophyll biosynthesis and breakdown by BCMs maintains chlorophyll homeostasis during leaf development. Nat Commun. 2020; 11: 1254.

