

Product no **AS09 431****Anti-Cytokinin | Cis-zeatin riboside (for immunolocalization)****Product information**

Immunogen	BSA-conjugated, via riboside, cis-zeatin riboside
Host	Rabbit
Clonality	Polyclonal
Purity	Total IgG. Protein G purified in PBS.
Format	Liquid
Quantity	200 µg
Storage	Store lyophilized/reconstituted at -20°C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please remember to spin the tubes briefly prior to opening them to avoid any losses that might occur from material adhering to the cap or sides of the tube.

Additional information | Cytokinin | Cis-zeatin riboside (for immunolocalization)**Application information**

Recommended dilution	Specific information about dilution is going to be included on the vial
Confirmed reactivity	Cis-zeatin riboside
Predicted reactivity	Cis-zeatin riboside
Not reactive in	No confirmed exceptions from predicted reactivity are currently known
Additional information	For 1 mg of antibodies For ELISA and immunoassay or For ELISA kit - please inquire, Antibodies are supplied in 50% glycerol, This antibody cannot discriminate between the zeatin riboside and zeatin, Zeatin cross-reactivity is usually lower ca,50%,BSA can be used For blocking using this antibody
Selected references	Han et al. (2019). Characterization and T-DNA insertion sites identification of a multiple-branches mutant br in <i>Betula platyphylla</i> and <i>Betula pendula</i> . <i>BMC Plant Biol.</i> 2019 Nov 12;19(1):491. doi: 10.1186/s12870-019-2098-y..