

Product no **AS09 431****Anti-Cytokinin | Cis-zeatin riboside (for immunolocalization)****Product information**

<b>Immunogen</b>	BSA-conjugated, via riboside, cis-zeatin riboside
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Purity</b>	Total IgG. Protein G purified in PBS.
<b>Format</b>	Liquid
<b>Quantity</b>	200 µg
<b>Storage</b>	Store lyophilized/reconstituted at -20°C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please remember to spin the tubes briefly prior to opening them to avoid any losses that might occur from material adhering to the cap or sides of the tube.

**Additional information** Cytokinin | Cis-zeatin riboside (for immunolocalization)**Application information**

<b>Recommended dilution</b>	Specific information about dilution is going to be included on the vial
<b>Confirmed reactivity</b>	Cis-zeatin riboside
<b>Predicted reactivity</b>	Cis-zeatin riboside
<b>Not reactive in</b>	No confirmed exceptions from predicted reactivity are currently known
<b>Additional information</b>	For 1 mg of antibodies For ELISA and immunoassay or For ELISA kit - please inquire, Antibodies are supplied in 50% glycerol, This antibody cannot discriminate between the zeatin riboside and zeatin, Zeatin cross-reactivity is usually lower ca,50%,BSA can be used For blocking using this antibody
<b>Selected references</b>	<a href="#">Han et al. (2019)</a> . Characterization and T-DNA insertion sites identification of a multiple-branches mutant br in Betula platyphylla and Betula pendula. BMC Plant Biol. 2019 Nov 12;19(1):491. doi: 10.1186/s12870-019-2098-y..