

Product no **AS09 443**

Anti-Cytokinin | Kinetin riboside (1 mg)

Product information

| | |
|-------------------------------|---|
| Immunogen | BSA-conjugated, via ribose, kinetin riboside |
| Host | Rabbit |
| Clonality | Polyclonal |
| Purity | Total IgG. Protein G purified in PBS. |
| Format | Lyophilized |
| Quantity | 1 mg |
| Reconstitution | For reconstitution add 50 µl of sterile water |
| Storage | Store lyophilized/reconstituted at -20°C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please remember to spin the tubes briefly prior to opening them to avoid any losses that might occur from material adhering to the cap or sides of the tube. |
| Additional information | Cytokinin Kinetin riboside (1 mg) |

Application information

| | |
|-----------------------------|--|
| Recommended dilution | 2 µl/15 ml |
| Confirmed reactivity | Kinetin riboside |
| Predicted reactivity | Kinetin riboside |
| Selected references | Bowie et al. (2018). N6-Furfuryladenine is protective in Huntington's disease models by signaling huntingtin phosphorylation. Proc Natl Acad Sci U S A. 2018 Jul 24;115(30):E7081-E7090. doi: 10.1073/pnas.1801772115. |