

Product no **AS09 439****Anti-Cytokinin | Meta-topolin riboside (5 mg)****Product information**

|                  |   |
|------------------|---|
| <b>Immunogen</b> | BSA-conjugated, via ribose, meta-topolin riboside   |
| <b>Host</b>      | Rabbit  |
| <b>Clonality</b> | Polyclonal  |
| <b>Purity</b>    | Total IgG. Protein G purified in PBS.   |
| <b>Format</b>    | Lyophilized   |
| <b>Quantity</b>  | 5 mg  |
| <b>Storage</b>   | Store lyophilized/reconstituted at -20°C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please remember to spin the tubes briefly prior to opening them to avoid any losses that might occur from material adhering to the cap or sides of the tube. |

**Additional information** | Cytokinin | Meta-topolin riboside (5 mg)**Application information**

|                             |  |
|-----------------------------|--|
| <b>Recommended dilution</b> | 2-10 µl/15 ml; Specific information about dilution is going to be included on the vial |
| <b>Confirmed reactivity</b> | Meta-topolin riboside  |
| <b>Predicted reactivity</b> | Meta-topolin riboside  |
| <b>Not reactive in</b>      | No confirmed exceptions from predicted reactivity are currently known                  |