


Product no **AS15 3082****Anti-DET1 | Regulator of the proteasomal degradation of LHY****Product information**

<b>Immunogen</b>	Recombinant, full length DET1 of <i>Arabidopsis thaliana</i> , overexpressed in <i>E.coli</i> , UniProt: <a href="#">P48732</a> , TAIR: <a href="#">AT4G10180</a>
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Purity</b>	Serum
<b>Format</b>	Lyophilized
<b>Quantity</b>	50 µl
<b>Reconstitution</b>	For reconstitution add 50 µl of sterile water
<b>Storage</b>	Store lyophilized/reconstituted at -20°C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please remember to spin the tubes briefly prior to opening them to avoid any losses that might occur from material adhering to the cap or sides of the tube.

**Additional information** | This product can be sold containing ProClin if requested

**Application information**

<b>Recommended dilution</b>	1 : 2500 (WB)
<b>Expected   apparent MW</b>	62   62 kDa
<b>Confirmed reactivity</b>	<i>Arabidopsis thaliana</i>
<b>Predicted reactivity</b>	<i>Capsicum annuum</i> , <i>Glycine max</i> , <i>Medicago truncatula</i> , <i>Oryza sativa</i> , <i>Populus trichocarpa</i> , <i>Ricinus communis</i> , <i>Solanum lycopersicum</i> , <i>Solanum tuberosum</i> , <i>Theobroma cacao</i> , <i>Zea mays</i> , <i>Zostera marina</i> Species of your interest not listed? <a href="#">Contact us</a>
<b>Not reactive in</b>	No confirmed exceptions from predicted reactivity are currently known
<b>Selected references</b>	<a href="#">Castells et al (2011)</a> . The conserved factor DE-ETIOLATED 1 cooperates with CUL4-DDB1DDB2 to maintain genome integrity upon UV stress. EMBO J. 2011 Mar 16;30(6):1162-72. doi: 10.1038/emboj.2011.20. Epub 2011 Feb 8.

**application example**

Recombinant protein (**0.5-1-1.5-2 µg of protein**) and total proteins from *Arabidopsis thaliana* whole leaves from plants grown in control light (**C**), low light (**LL**) and high light (**HL**), corresponding to 1 µg of chlorophylls, were extracted with loading buffer (10% glycerol, 62.5 mM Tris pH 6.8, 2% SDS, 5% β-mercaptoethanol) and denatured at 100°C (boiling water) for 1 min. Proteins were separated on 12% SDS-PAGE (Laemly) and blotted 1h to PVDF using tank transfer. Blots were blocked with blocking solution (PBS 1X, 0.2% w/v Tween, 5% powder milk) for 1h at room temperature (RT) with agitation. Blot was incubated in the primary antibody diluted in blocking solution, at a dilution of 1 : 25,000 for 1h at RT with agitation. The antibody solution was decanted and the blot was rinsed briefly twice, then washed 3 times for 10 min in blocking solution at RT with agitation. Blot was incubated in secondary antibody (anti-rabbit IgG alkaline phosphatase conjugated) diluted to 1:30 000 in blocking buffer for 1h at RT with agitation. The blot was washed 2 times for 10 min in blocking solution and once with PBS 1X solution for 10 min, then developed in developing buffer developing buffer NBT/BCIP by manual agitation.

Courtesy of Stefano Cazzaniga, University of Verona, Italy