

Product no **AS10 1019****Donkey anti-Rabbit IgG (H&L), TRITC conjugated, min, cross-reactivity to bovine, chicken, goat, guinea pig, hamster, horse, human, mouse, rat, sheep IgG****Product information****Host** | Donkey**Clonality** | Polyclonal**Purity** | Immunogen affinity purified IgG.**Format** | Lyophilized**Quantity** | 1 mg**Reconstitution** | For reconstitution add 1.1 ml of sterile water. Let it stand 30 minutes at room temperature to dissolve. Prepare fresh working dilutions daily**Storage** | Store lyophilized material at 2-8°C. For long time storage after reconstitution, dilute the antibody solution with glycerol to a final concentration of 50% glycerol and store as liquid at -20°C, to prevent loss of enzymatic activity. For example, if you have reconstituted 1 mg of antibody in 1.1 ml of sterile water add 1.1 ml of glycerol. Such solution will not freeze in -20°C. If you are using a 1:5000 dilution prior to diluting with glycerol, then you would need to use a 1:2500 dilution after adding glycerol. Prepare working dilution prior to use and then discard. Be sure to mix well but without foaming.**Additional information** | This antibody reacts with the heavy chains on rabbit IgG and with the light chains on all rabbit immunoglobulins based on immunoelectrophoresis.

Minimum cross-reactivity is observed to non-immunoglobulin rabbit serum proteins based on immunoelectrophoresis.

Antibody is supplied in 10 mM sodium phosphate, 150 mM sodium chloride, pH 7.2, 1% (w/v) BSA, protease/IgG free and 0.05 % (w/v) sodium azide as preservative.

Application information**Recommended dilution** | The optimal working dilution should be determined by the investigator, (ICC), immunohistochemistry (IHC) (Flow cyt)