


Product no **AS10 935****Dsp | Death-specific protein****Product information**

Immunogen	KLH-conjugated synthetic peptide derived from known sequences of DSP in diatoms, conserved in Dsp1 and Dsp2, UniProt: B6RD07 of <i>Thalassiosira pseudonana</i> and <i>Skeletonema costatum</i> and Dsp1 in <i>Fragilariopsis cylindrus</i> and <i>Phaeodactylum tricornutum</i>
Host	Rabbit
Clonality	Polyclonal
Purity	Serum
Format	Lyophilized
Quantity	200 µl
Reconstitution	For reconstitution add 200 µl of sterile water
Storage	Store lyophilized/reconstituted at -20°C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please remember to spin the tubes briefly prior to opening them to avoid any losses that might occur from material adhering to the cap or sides of the tube.

Application information

Recommended dilution	1 : 10 000 (WB)
Expected apparent MW	25 25 kDa
Confirmed reactivity	<i>Emiliana huxleyi</i> , <i>Aureococcus anophagefferens</i> , <i>Fragilariopsis cylindrus</i> , <i>Phaeodactylum tricornutum</i> , <i>Skeletonema costatum</i> , <i>Thalassiosira pseudonana</i>
Predicted reactivity	Species of your interest not listed? Contact us
Not reactive in	<i>Aureococcus anophagefferens</i>
Selected references	Thamatrakoln et al. (2013) . Death-specific protein in a marine diatom regulates photosynthetic responses to iron and light availability. PNAS 2013 Dec 10;110(50):20123-8. doi: 10.1073/pnas.1304727110. Epub 2013 Nov 25.

Application example

5 µg of total protein, extracted with LDS sample buffer (PSB) were separated on 4-20% SDS-PAGE and transferred to PVDF. Blot was blocked with 2% ECL block in TBS-T (0.1% Tween-20) for 1h at room temperature (RT) with agitation. Blot was incubated in primary antibody at a dilution of 1: 10,000 for 1h at RT with agitation. Antibody solution was decanted and the blot rinsed briefly twice with TBS-T, then washed once for 15 min and 3 times for 5 min. at RT with agitation. Blot was incubated in secondary antibody (horse radish peroxidase conjugated anti-rabbit IgG) at a dilution of 1:20 000, for 1h at RT with agitation. The blot was washed as above and developed for 5 min with ECL according to the manufacturer's instructions. Image (20 sec. exposure) captured with a BioRad Chemidoc MP.

Predicted activity and molecular weights (Da) - *Thalassiosira pseudonana* Dsp1 (24572 Da) Dsp2 (25406 Da), *Skeletonema costatum* Dsp1 (24848 Da) Dsp2 (26566 Da), *Phaeodactylum tricornutum* Dsp1 (26585 Da, *Fragilariopsis cylindrus* Dsp1 (20312 Da).

Competition assay (right half of blot): Same protein loads as for lanes 1-7. Antisera were pre-incubated for 90 min. in TBS-T with of an excess of the free peptide antigen

Courtesy of Dr. K. Thamatrakoln unpublished data; rights retained by the contributor.