


Product no **AS14 2782****Anti-EPSP synthase | 3-phosphoshikimate 1-carboxyvinyltransferase (chloroplastic, *Arabidopsis thaliana*)****Product information**

Immunogen	KLH-conjugated peptide, derived from <i>Arabidopsis thaliana</i> EPSP synthase, UniProt: P05466 TAIR: At2g45300
Host	Rabbit
Clonality	Polyclonal
Purity	Serum
Format	Lyophilized
Quantity	50 µl
Reconstitution	For reconstitution add 50 µl of sterile water
Storage	Store lyophilized/reconstituted at -20 °C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please remember to spin the tubes briefly prior to opening them to avoid any losses that might occur from material adhering to the cap or sides of the tube.

Application information


Recommended dilution	1 : 1000 (WB)
Expected apparent MW	55 kDa
Confirmed reactivity	<i>Arabidopsis thaliana</i>
Predicted reactivity	<i>Arachis hypogaea</i> , <i>Cannabis sativa</i> , <i>Coffea arabica</i> , <i>Cucumis sativus</i> , <i>Glycine max</i> , <i>Glycine soja</i> , <i>Malus domestica</i> , <i>Nicotiana tabacum</i> , <i>Zea mays</i> , <i>Vitis vinifera</i> Species of your interest not listed? Contact us

application example

0.5 µg of purified *Arabidopsis thaliana* EPSPS was separated on 10 % SDS-PAGE and blotted 2h to nitrocellulose Hibond ECLmembrane (GE Healthcare) using semi-dry or tank transfer. Blots were blocked with for 1h at room temperature (RT) with agitation. Blot was incubated in the primary antibody in TBS-T with (0.1 % Tween 20) at a dilution of 1: 1 000 for 1h at RT with agitation. The antibody solution was decanted and the blot was rinsed briefly twice, then washed once for 15 min and 3 2 times for 5 min in TBS-T at RT with agitation. Blot was incubated in secondary antibody (anti-rabbit IgG horse radish peroxidase conjugated, from Agrisera [AS09 602](#)) diluted to 1: 20 000 in for 1h at RT with agitation. The blot was washed as above and developed for 5 min with ECL according to the manufacturer's instructions. Protein detection was done by Image Quant LAS4000 (GE-Health Care) digital imager system with exposure time was of 5 minutes.

Recombinant *Arabidopsis thaliana* EPSPS protein (aka 3-phosphoshikimate 1-carboxyvinyltransferase) used in the assay was identified by LCMS.

Courtesy of Dr. Altanbadral Sharkhuu, King Abdullah University of Science and Technology, Thuwal, Saudi Arabia



50 µg of total protein from shoots of 18 d old *Arabidopsis thaliana* were extracted with buffer (125 mM Tris-HCl pH6.8; 12% SDS; 10% glycerol, 22% b-mercaptoethanol; 0.001% bromophenol blue) were separated on % SDS-PAGE and blotted 3h to nitrocellulose membrane using tank transfer. Blots were blocked with TBST with 5% skim milk for 1h at room temperature (RT) with agitation. Blot was incubated in the primary antibody (Rabbit-EPSPS) at a dilution of 1: 1 000 for over night at 4 C with agitation. The antibody solution was decanted and the blot was rinsed briefly twice, then washed once for 15 min and 3 times for 5 min in TBS-T at RT with agitation. Blot was incubated in secondary antibody (anti-rabbit IgG horse radish peroxidase conjugated, from Agrisera [AS09 602](#)) diluted to 1:20 000 in for 1h at RT with agitation. The blot was washed as above and developed for 5 min with ECL according to the manufacturer's instructions. Detection was done by Image Quant LAS 4000 (GE-Health Care). Exposure time was 2 min.



This product is **for research use only** (not for diagnostic or therapeutic use)

contact: support@agrisera.com

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | www.agrisera.com

Courtesy of Dr. Altanbadralt Sharkhuu, King Abdullah University of Science and Technology, Thuwal, Saudi Arabia