


Product no **AS20 4436****FNR | Ferredoxin NADP Reductase (root)****Product information**

Immunogen	Purified full length, tag cleaved, recombinant maize root FNR (R-FNR), UniProt: Q41736
Host	Rabbit
Clonality	Polyclonal
Purity	Total IgG. Protein A purified in PBS, 50% glycerol. Filter sterilized.
Format	Liquid at 1 mg/ml.
Quantity	100 µg
Storage	Store at -20°C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please remember to spin the tubes briefly prior to opening them to avoid any losses that might occur from material adhering to the cap or sides of the tube.

Application information


Recommended dilution	1: 1000 - 1: 30 000 (WB)
Expected apparent MW	36,3 kDa 35,57 kDa
Confirmed reactivity	<i>Arabidopsis thaliana</i> , <i>Triticum aestivum</i> , <i>Secale cereale</i> , <i>Zea mays</i>
Predicted reactivity	Species of your interest not listed? Contact us
Not reactive in	No confirmed exceptions from predicted reactivity are currently known
Additional information	Antibody will recognize two root FNR proteins (R-FNR1 and R-FNR2) of <i>Arabidopsis thaliana</i> and <i>Zea mays</i> as well as leaf FNRs. This antibody can be used as a marker of plastids localised in roots.
Selected references	Hachiya et al. (2016) . Arabidopsis Root-Type Ferredoxin:NADP(H) Oxidoreductase 2 Is Involved in Detoxification of Nitrite in Roots . Plant Cell Physiol. 57(11):2440-2450. doi: 10.1093/pcp/pcw158. Hachiya et al. (2016) . Arabidopsis Root-Type Ferredoxin:NADP(H) Oxidoreductase 2 Is Involved in Detoxification of Nitrite in Roots. Plant Cell Physiol. 57(11):2440-2450. doi: 10.1093/pcp/pcw158. Onda et al. (2000) . Differential Interaction of Maize Root ferredoxin:NADP(+) Oxidoreductase With Photosynthetic and Non-Photosynthetic Ferredoxin Isoproteins. Plant Physiol. 123(3):1037-45. doi: 10.1104/pp.123.3.1037. Onda et al. (2000) . Differential Interaction of Maize Root ferredoxin:NADP(+) Oxidoreductase With Photosynthetic and Non-Photosynthetic Ferredoxin Isoproteins. Plant Physiol. 123(3):1037-45. doi: 10.1104/pp.123.3.1037.



Secale cereale total chloroplast extract (leaves) (1), *Triticum aestivum* total chloroplast extract (leaves) (2), *Triticum aestivum* cytosolic protein extract (roots) (3), *Secale cereale* cytosolic protein extract (roots) (4), *Triticum aestivum* total protein extract (leaves) (5), *Secale cereale* total protein extract (leaves) (6), *Secale cereale* total plastid extract (roots) (7), *Secale cereale* total protein extract (roots) (8)

15 µg of protein extracted from leaves and roots of *Triticum aestivum* and *Secale cereale* were separated on 10% SDS-PAGE and blotted 1h onto nitrocellulose (0.45µm) using semi-dry transfer. After blocking with 5% milk in TBST, blots were incubated with the primary antibody at a dilution of 1:5000 in TBST for 1h at room temperature. Following incubation and wash steps, blots were incubated with secondary Goat anti-Rabbit IgG (H&L), HRP conjugated ([AS09 602](#), Agrisera) for 1 hour at a dilution of 1:40 000. Blots were developed with the HRP detection system using [AgriseraECLSuperBright](#).

Courtesy of Dr Bartosz Szabala, Department of Plant Genetics, Breeding and Biotechnology, Warsaw University of Life Sciences, Poland



10 µg/well of root total protein of *Arabidopsis thaliana* wild type leaf **(1)**, **(3)** and mutants *rfr1* **(2)** and *rfr2-2* **(4)** grown under 0.2 mM nitrate for 7 days were freshly extracted with 2x SDS-sample buffer (+ 2ME) for SDS-PAGE and denatured with 4X SDS buffer at 95°C for 5 min. Samples were separated on 10% SDS-PAGE and blotted 1h to PVDF membrane. Blot was blocked with 3 % skim milk/TBS-T, 1h/RT with agitation. Blot was incubated in the primary antibody at a dilution of 1: 2000 in TBS-T for 1-2h/RT. The antibody solution was decanted and the blot was washed 4 times for 10 min in TBS-T at RT with agitation. Blot was incubated in matching secondary antibody (anti-rabbit IgG horse radish peroxidase conjugated) diluted to 1:10 000 in for 1h/RT with agitation. The blot was washed as above and developed with a chemiluminescent detection reagent, following manufacture's recommendation.

R-FNR2 is a dominant form in wild type roots.

Mutant *rfr1* produces R-FNR2 and mutant *rfr2* produces R-FNR1.