


Product no **AS10 689****Anti-GluTR | Glutamyl -tRNA reductase****Product information****Immunogen** | KLH-conjugated peptide derived from available glutamyl-tRNA reductase sequences including *Arabidopsis thaliana* P49294**Host** | Rabbit**Clonality** | Polyclonal**Purity** | Serum**Format** | Lyophilized**Quantity** | 200 µl**Reconstitution** | For reconstitution add 200 µl of sterile water**Storage** | Store lyophilized/reconstituted at -20°C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please remember to spin the tubes briefly prior to opening them to avoid any losses that might occur from material adhering to the cap or sides of the tube.**Additional information** | Antibody reacts with recombinant GluTR isoforms: AtGluTR1, AtGluTR2 and NtGluTR1 (At - *Arabidopsis thaliana*, Nt - *Nicotiana tabacum*).**Application information****Recommended dilution** | 1 : 5000 (WB)**Expected | apparent MW** | 58 | kDa (*Arabidopsis thaliana*)**Confirmed reactivity** | *Arabidopsis thaliana*, *Hordeum vulgare*, *Nicotiana tabacum***Predicted reactivity** | *Brassica napus*, *Chlamydomonas reinhardtii*, *Cucumis sativus*, *Glycine max*, *Hordeum vulgare*, *Oryza sativa*, *Physcomitrium patens*, *Picea sitchensis*, *Pisum sativum*, *Populus trichocarpa*, *Solanum tuberosum*, *Sorghum bicolor*, *Ricinus communis*, *Zea mays*, *Vitis vinifera*Species of your interest not listed? [Contact us](#)**Not reactive in** | Cyanobacteria**Selected references** | Pavlovič et al. (2024). Diethyl ether anaesthesia inhibits de-etiolation of barley seedlings by locking them in intermediate skoto-photomorphogenetic state. *Physiol Plantarum*, Volume176, Issue 1.
Agrawal et al. (2020). The Functions of Chloroplast Glutamyl-tRNA in Translation and Tetrapyrrole Biosynthesis. *Plant Physiol.* 2020 Feb 18. pii: pp.00009.2020. doi: 10.1104/pp.20.00009
Montandon et al. (2019). In vivo trapping of proteins interacting with the chloroplast CLPC1 chaperone; potential substrates and adaptors. *J Proteome Res.* 2019 May 9. doi: 10.1021/acs.jproteome.9b00112.
Nishimura et al. (2013). ClpS1 Is a Conserved Substrate Selector for the Chloroplast Clp Protease System in *Arabidopsis*. *The Plant Cell* June 2013.**application example**

10 µl of leaf extract which was equivalent to 1 mg leaf material was loaded per lane, which may also correspond to approximately 50 µg protein. *Arabidopsis thaliana* seedlings were grown on vermiculite for 3 weeks under continuous illumination at a light intensity of 80 µE m⁻² s⁻¹ at 22°C. Twenty mg leaf material was collected from mature leaves and extracted with 200 µl of the tissue homogenization buffer. 10 µl of leaf extract which was equivalent to 1 mg leaf material was loaded per lane. Detection Protocol: Leaf protein was separated on 14% SDS-PAGE and blotted 2h to PVDF membrane from GE Healthcare. The blot was blocked with PBS-T (PBS plus 0.1% tween 20) containing 3% skim milk for 1h at room temperature (RT: approximately 22 degrees C) with gentle agitation. The blots were briefly washed twice with PBS-T and then incubated with anti-GluTR antibody which was diluted 1:1000 with PBS-T for 1h at RT with agitation. The primary antibody solution was decanted and the blot was rinsed twice, when washed once for 10 min and 3 times for 5 min in PBS-T containing 0.5% (w/v) skim milk at RT with agitation. The blot was incubated with the secondary antibody (HRP-conjugated anti-rabbit IgG) which was diluted 1:20 000 with PBS-T containing 0.5% (w/v) skim milk for 1h at RT with agitation. The blot was washed as described above and incubated with Western Lightning Plus-ECL from Perkin-Elmer for 1 min. The chemiluminescent signal was captured with a CCD camera (LumiVision: Aisin Seiki Inc. Aichi, Japan) for 60 s.

Plant material grown for the experiment illustrated below: barley seeds were sown on vermiculite and grown for 6 days in darkness at 22°C. Subsequently, seedlings were illuminated for 24 hours under a light intensity of 80 $\mu\text{E m}^{-2} \text{s}^{-1}$. Approximately, 100 mg of *Hordeum vulgare* leaf material was harvested before illumination **(1)**, and 2 h **(2)**, 6 h **(3)** and 24 hours **(4)** after the onset of light; *Arabidopsis thaliana* wt **(5)**. Each leaf material was extracted with 1 ml of the tissue homogenization solution (50 mM Tris Cl pH8.0, 2% Lithium Dodesyl Sulfate, 12% sucrose, 1.5% dithiothreitol).

Courtesy of Kaori TAKAHASHI and Ryouichi TANAKA (Hokkaido University)