

This product is for research use only (not for diagnostic or therapeutic use)

contact: support@agrisera.com

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | www.agrisera.com

Product no AS11 1823

Goat anti-Chicken IgY (H&L), DyLight® 594 conjugated

Product information

Immunogen purified chicken IgY, whole molecule

Host Goat

Clonality Polyclonal

Purity Immunogen affinity purified goat IgG.

Format Lyophilized

Quantity 1 mg

Reconstitution

For reconstitution add 1 ml of sterile water, Allow reconstituted product to stand for at least 30 minutes at room temperature prior to dilution, If necessary, centrifuge to remove any particulates, Prepare fresh working dilution daily

Storage

Store lyophilized material at 2-8°C. Product is stable for 4 weeks at 2-8°C after rehydration. For long time storage after reconstitution, dilute the antibody solution with glycerol to a final concentration of 50% glycerol and store as liquid at -20°C, to prevent loss of enzymatic activity. For example, if you have reconstituted 1 mg of antibody in 1,1 ml of sterile water add 1,1 ml of glycerol. Such solution will not freeze in -20°C, If you are using a 1:5000 dilution prior to diluting with glycerol, then you would need to use a 1:2500 dilution after adding glycerol. Prepare working dilution prior to use and then discard. Be sure to mix well but without foaming.

Additional information

Conjugate is present in 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Protease/lgG free. 0.05 % (w/v) sodium azide is added as preservative.

DyLight® 594 (Ex = 593 nm; Em = 618 nm).

Application information

Recommended dilution 1:50-1:5 000 (ICC), 1:20 -1:2000 (IHC)

Confirmed reactivity Chicken IgY heavy and light chains (H&L)

Predicted reactivity Chicken IgY Heavy and Light chains (H&L)

Not reactive in No confirmed exceptions from predicted reactivity are currently known

Additional information | Based

Based on immunoelectrophoresis this antibody reacts with heavy chains on chicken IgG and light chains on all chicken immunoglobulins.

No reactivity is observed to non-immunoglobulin chicken serum proteins based in immunoelectrophoresis.