

Goat anti-Human IgG F(ab')₂ Affinity purified, Unconjugated

Qty: AS21 4574



AS21 4574 | Clonality: **Polyclonal** | Host: **Goat** | Reactivity: **Human IgG F(ab')₂ with no reactivity to:**

Fc fragment of human IgG
non-immunoglobulin human serum proteins
serum proteins from: bovine, horse or mouse

Price: 218 €

Antibody protocols

- Product Info
- Immunogen: Purified human IgG F(ab')₂ fragment
- Host: Goat
- Clonality: Polyclonal
- Purity: Immunogen affinity purified goat IgG.
- Format: Liquid
- Quantity: 2 mg
- Storage: Store at 2-8°C. Shelf life is one year from date of receipt.
- Tested applications: ELISA (ELISA) Immunoprecipitation (IP), Immunohistochemistry (IHC)
- Recommended dilution: The optimal working dilution should be determined by the investigator
- Reactivity
- Confirmed reactivity: F(ab')₂ of human IgG
- Predicted reactivity: F(ab')₂ of human IgG
- Application Examples



Goat anti-human IgG F(ab')₂ antibodies were separated on SDS-PAGE. Following the staining, two predominant bands at 50 and 24 kDa are observed.

- Additional Information
- Antibody is supplied in 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 0.05 % (w/v) of sodium azide as preservative. It is a clear, colourless liquid, filter sterilized.

Additional information:

Antibody purity ≥95% based on SDS-PAGE.
 Based on IEP this antibody reacts with: F(ab')₂ fragment of human IgG.

Additional information
 (application):

Based on IEP no reactivity is observed to:
 Fc fragment of human IgG
 non-immunoglobulin human serum proteins
 serum proteins from bovine, horse and mouse

- Background
- Background: **Goat anti-human IgG (H&L), F(ab)'2 fragment** is a secondary antibody which binds to human IgG (H&L), F(ab)'2 fragment in [immunological assays](#).