

This product is for research use only (not for diagnostic or therapeutic use)

contact: support@agrisera.com

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | www.agrisera.com

## Product no AS16 3354

Goat anti-Human IgG Fc, F(ab)'2 Fragment, FITC conjugated, min, cross-reactivity to bovine, horse, mouse or rabbit serum proteins, mouse IgG1

## **Product information**

Immunogen Purified Human IgG, Fc fragment

**Host** Goat

Clonality Polyclonal

**Purity** Immunogen affinity purified goat IgG.

Format Liquid

Quantity 0,5 mg

Storage Store at 2-8°C.

**Additional information** 

Antibody purity is > 90% based on SDS-PAGE Small amounts of intact IgG may be present.

Affinity purified using solid phase Human IgG .

 $Antibody \ is \ supplied \ in \ 10 \ mM \ Sodium \ Phosphate, \ 0.15 \ M \ Sodium \ Chloride, \ pH \ 7.2, \ 1 \ \% \ (w/v) \ BSA, \ Protease/lgG \ free.$ 

0.05% (w/v) Sodium Azide is added as a preservative.

Based on IEP, this antibody reacts with: • heavy ( ) chains on human IgG .

Based on IEP, no reactivity is observed to: • non-immunoglobulin human serum immunoglobulins • light chains on all human immunoglobulins • human IgG F(ab)'2 fragment • mouse IgG1 • serum proteins from bovine, horse, mouse, or

## **Application information**

Recommended dilution 1:20-1:2000 (Flow cyt), (IL)

Confirmed reactivity Goat anti-Human IgG Fc, F(ab)'2 Fragment