

Product no **AS16 3358**

Goat anti-Human IgG Fc, F(ab)'2 Fragment, TRITC conjugated, min, cross-reactivity to bovine, horse, mouse or rabbit serum proteins, mouse IgG1

Product information

| | |
|-----------------------|---|
| Immunogen | Purified Human IgG, Fc fragment |
| Host | Goat |
| Clonality | Polyclonal |
| Purity | Immunogen affinity purified goat IgG. |
| Format | Lyophilized |
| Quantity | 0.5 mg |
| Reconstitution | For reconstitution add 0.55 ml of sterile water |
| Storage | Store at 2-8 °C. |

| | |
|-------------------------------|---|
| Additional information | <p>Antibody purity is > 90% based on SDS-PAGE. Small amounts of intact IgG may be present. Affinity purified using solid phase Human IgG.</p> <p>Antibody is supplied in 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1% (w/v) BSA, Protease/IgG free. 0.05% (w/v) Sodium Azide is added as a preservative.</p> <p>Based on IEP, this antibody reacts with: • heavy () chains on human IgG.</p> <p>Based on IEP, no reactivity is observed to: • non-immunoglobulin human serum immunoglobulins • light chains on all human immunoglobulins • human IgG F(ab)'2 fragment • mouse IgG1 • serum proteins from bovine, horse, mouse, or rabbit.</p> |
|-------------------------------|---|

Application information

| | |
|-----------------------------|--|
| Recommended dilution | 1 : 20-1 : 2000 (Flow cyt), (IL) |
| Confirmed reactivity | Goat anti-Human IgG Fc, F(ab)'2 Fragment |