

Product no **AS16 3349****Goat anti-Human IgG Fc, TRITC conjugated, min, cross-reactivity to human IgA + IgM, bovine, horse, mouse or rabbit serum proteins****Product information****Immunogen** Purified Human IgG, Fc fragment**Host** Goat**Clonality** Polyclonal**Purity** Immunogen affinity purified goat IgG.**Format** Lyophilized**Quantity** 1 mg**Reconstitution** For reconstitution add 1,1 ml of sterile water, Let it stand 30 minutes at room temperature to dissolve, Prepare fresh working dilutions daily**Storage** Store at 2-8°C.**Additional information** Antibody purity is > 95% based on SDS-PAGE.
Affinity purified using solid phase Human IgG .
Antibody is supplied in 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Protease/IgG free.
0.05% (w/v) Sodium Azide is added as a preservative.Based on IEP, this antibody reacts with: • heavy (γ) chains on human IgG .

Based on IEP, no reactivity is observed to: • non-immunoglobulin human serum immunoglobulins • light chains on all human immunoglobulins • human IgA or IgM • serum proteins from bovine, mouse, or rabbit .

Application information**Recommended dilution** 1 : 20-1 : 2000 (Flow cyt), (IL)**Confirmed reactivity** Human IgG Fc