

This product is for research use only (not for diagnostic or therapeutic use)

contact: support@agrisera.com

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | www.agrisera.com

Product no AS10 1237

Goat anti-Human IgM (µ chain), F(ab)'2 fragment, ALP conjugated

Product information

Host Goat

Clonality Polyclonal

Purity Immunogen affinity purified IgG, F(ab)'2 fragment.

Format Liquid

Quantity 0.5 mg

Storage

Non-diluted antibody is stable for 4 years at 2-8 °C. For storage at -20 °C dilute antibody solution with an equal volume of glycerol to obtain final glycerol concentration of 50% to prevent loss of enzymatic activity. Such solution will not freeze in -20 °C. If you are using a 1:5000 dilution prior to diluting with glycerol, then you would need to use a 1:2500 dilution after adding glycerol. Prepare working dilution prior to use and then discard, Be sure to mix well but without foaming.

Additional information

This antibody reacts with:heavy (µ) chains on human IgM based on immunoelectrophoresis.

Minimum cross-reactivity is observed to: non-immunoglobulin serum proteins human IgG.

Antibody is supplied in 30 mM triethanolamine, pH 7.2, 5 mM magnesium chloride, 0.1 mM zinc chloride, 1% (w/v) BSA, Protease/IgG free, 0.05% (w/v) sodium azide as preservative.

Concentration: 1.50 mg/ml (E 1% at 280 nm = 13.0)

Application information

Recommended dilution 1:50-1:5000 (ICC), 1:20-1:2000 (IHC), 1:500-1:2000 (WB)