

Product no **AS10 1298****Goat anti-Human IgM (μ chain), F(ab)'2 fragment, ALP conjugated, min, cross-reactivity to human IgG or IgA****Product information**

Host	Goat
Clonality	Polyclonal
Purity	Immunogen affinity purified IgG, F(ab)'2 fragment.
Format	Liquid
Quantity	0.5 mg
Storage	Non-diluted antibody is stable for 4 years at 2-8°C. For storage at -20°C dilute antibody solution with an equal volume of glycerol to obtain final glycerol concentration of 50 % to prevent loss of enzymatic activity. Such solution will not freeze in -20°C. If you are using a 1:5000 dilution prior to diluting with glycerol, then you would need to use a 1:2500 dilution after adding glycerol. Prepare working dilution prior to use and then discard. Be sure to mix well but without foaming.

Additional information	This antibody reacts with:heavy (μ) chains on human IgM based on immunoelectrophoresis. Minimum cross-reactivity is observed to: non-immunoglobulin serum proteins human IgG or IgA. Antibody is supplied in 30 mM triethanolamine, pH 7.2, 5 mM magnesium chloride, 0.1 mM zinc chloride, 1% (w/v) BSA, Protease/IgG free, 0.05 % (w/v) sodium azide as preservative.
-------------------------------	--

Application information**Recommended dilution** | 1 : 50-1 : 5000 (ICC) , 1 : 20-1 : 2000 (IHC), (Flow cyt), 1 : 500-1 : 2000 (WB)