

Product no **AS16 3441****Goat anti-Mouse IgG (H&L), F(ab)'2 fragment, ALP conjugated, min, cross-reactivity to bovine, goat, human, rabbit, or rat IgG****Product information**

| | |
|------------------|---------------------------------------|
| Immunogen | Purified Mouse IgG, whole molecule |
| Host | Goat |
| Clonality | Polyclonal |
| Purity | Immunogen affinity purified goat IgG. |
| Format | Liquid |
| Quantity | 0.5 mg |
| Storage | Store at 2-8°C. |

| | |
|-------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Additional information | Antibody purity is > 90% based on SDS-PAGE. Small amounts of intact IgG may be present. Affinity purified using solid phase Mouse IgG. Antibody is supplied in 30 mM Triethanolamine, pH 7.2, 5 mM Magnesium Chloride, 0.1 mM Zinc Chloride, 1% (w/v) BSA, Protease/IgG free. Based on IEP, this antibody reacts with: • heavy () chains on mouse IgG • light chains on all mouse immunoglobulins. Based on IEP, no reactivity is observed to: • non-immunoglobulin human serum immunoglobulins • bovine, goat, human, rabbit or rat IgG. |
|-------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|

Application information

| | |
|-----------------------------|-----------------------------------|
| Recommended dilution | 1 : 20-1 : 2000 (Flow cyt), (IL) |
| Confirmed reactivity | Mouse IgG (H&L), F(ab)'2 Fragment |