

This product is **for research use only** (not for diagnostic or therapeutic use)

contact: support@agrisera.com

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | www.agrisera.com

## Product no AS10 1017

## Goat anti-Rabbit IgG (H&L), F(ab)'2 fragment, ALP (alkaline phosphatase) conjugated, min, cross-reactivity to bovine,human, mouse IgG/serum

## **Product information**

Host	Goat
Clonality	Polyclonal
Purity	Immunogen affinity purified goat IgG, F(ab)'2 fragment.
Format	Liquid
Quantity	0.5 mg
Ĵ	Store non-diluted antibody at 2-8°C. For storage at -20°C dilute antibody solution with an equal volume of glycerol to obtain final glycerol concentration of 50 % to prevent loss of enzymatic activity. Such solution will not freeze in -20°C. If you are using a 1:5000 dilution prior to diluting with glycerol, then you would need to use a 1:2500 dilution after adding glycerol. Prepare working dilution prior to use and then discard, Be sure to mix well but without foaming.
Additional information	immunoelectrophoresis. Minimum cross-reactivity is observed to: non-immunoglobulin rabbit serum proteins IgG from bovine, goat, human, mouse or rat.
	Antibody is supplied in 30 mM triethanolamine, pH 7.2, 5 mM magnesium chloride, 0.1 mM zinc chloride, 1 % (w/v) BSA, Protease/IgG free and , 0.05 % (w/v) sodium azide as preservative.
Application information	

Recommended dilution 1 : 50-1 : 5000 (ICC), 1 : 20-1 : 2000 (IHC), 1 : 500-1 : 2000 (WB)