



## Goat anti-Rat IgM Fc, Biotin conjugated

Qty: AS25 5117

**AS25 5117** | Clonality: **Polyclonal** | Host: **Goat** | Reactivity: **Rat IgM (Fc)**

Price: 330 €

- Product Info
- Immunogen: Purified rat IgM
- Host: Goat
- Clonality: Polyclonal
- Format: Lyophilized
- Quantity: 1 ml reconstituted
- Reconstitution: For reconstitution, add 1 ml of sterile water. Let it stand 30 minutes at room temperature to dissolve. Spin centrifuge shortly to remove any particles. Prepare fresh working dilutions daily by adding sterile phosphate buffered saline (PBS, pH 7.2).
- Storage: Store lyophilized material at 2-8°C. For long time storage at or below -20°C; once reconstituted, make aliquots to avoid repeated freeze-thaw cycles. Please, remember to spin tubes briefly prior to opening them to avoid any losses that might occur from lyophilized material adhering to the cap or sides of the tubes.
- Tested applications: ELISA /ELISA), Immunocytochemistry (ICC), Immunohistochemistry (IHC) (paraffin), Dot blot (Dot), Western blot (WB)
- Recommended dilution: The optimum working dilution of each conjugate should be established by titration before being used.
- Reactivity
- Confirmed reactivity: Fc part of the rat IgM
- Not reactive in: Rat IgG, IgA and IgG/Fab or any non-Ig protein in rat serum.
- Additional Information
- Additional information: Biotin conjugated goat IgG in PBS pH 7.2. No preservative is added.
- Background
- Background: **Goat anti-rat IgM (Fc)** is a secondary antibody conjugated to biotin which binds to the Fc region of the rat IgM.