

This product is for research use only (not for diagnostic or therapeutic use)

contact: support@agrisera.com

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | www.agrisera.com

## Product no AS20 4515

## GAL | Beta galactosidase

## **Product information**

Immunogen Full length Beta galactosidase from E.coli UniProt: B7UJI9

**Host** Rabbit

Clonality Polyclonal

**Purity** Total IgG. Protein A purified in PBS, 50 % glycerol, filter sterilized.

Format Liquid at 2 mg/ml.

Quantity 200 μg

Storage

Store at -20°C, Avoid repeated freeze-thaw cycles. Please remember to spin the tubes briefly prior to opening them to avoid any losses that might occur from material adhering to the cap or sides of the tube. , Do not store this antibody


## **Application information**

Recommended dilution 1: 2000 - 1: 3000 (ELISA), 1: 200 - 1: 500 (IF, IP), 1: 1000 - 1: 2000 (WB)

Expected | apparent 116 kDa

**Confirmed reactivity** Beta galactosidase (*E.coli*) and beta galactosidase tagged proteins


Selected references



- 1. Purified -galactosidase, 10 ng
- 2. Uninduced E. coli K12 cell extract (30 ug)
- 3. E.coli K12 cell extract induced by IPTG for -galactosidase expression (30 ug)

The anti- -galactosidase antibody was used at 1:1000 dilution in TBS-T.

Molecular mass of -galactosidase is 116 kDa.



Sample: Crude extract of E. coli K12 cells induced by IPTG for -galactosidase expression.

- 1. Crude cell extract
- 2. Supernatant of the immuno-precipitated E. coli crude extract
- 3. Immuno-precipitate of crude E. coli extract


The anti- -galactosidase antibody was used at 1:500 dilution for immune-precipitation and 1:1000 for western blot. IgG (H) is heavy chain of IgG immunoglobulin. Membrane for transfer: nitrocellulose. Detection: chemiluminescent, following manufacture's recommendations.



This product is for research use only (not for diagnostic or therapeutic use)

contact: support@agrisera.com

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | www.agrisera.com



HEK293A cells were transfected with -Gal cDNA, fixed with 4% paraformaldehyde 24 hrs later, permeabilized with methanol, and immunostained with anti- -Gal antibody (1:500) and Alexa 555-conjugated rabbit IgG (1:500). Chromosomal DNA was stained with Hoechst 33342. Scale bar, 50 mm. Note that the antibody reacts only in transfected cells.