

This product is **for research use only** (not for diagnostic or therapeutic use)

contact: support@agrisera.com

Agrisera AB | Box 57 | SE-91112 Vännäs | Sweden | +46 (0)935 33 000 | www.agrisera.com

Product no **AS07 260-DL488**

## Anti-H<sub>+</sub>ATPase | Plasma membrane H<sub>+</sub>ATPase (rabbit antibody), DyLight® 488 conjugated (40 µg)

### Product information

<b>Immunogen</b>	KLH-conjugated synthetic peptide, derived from available di and monocot, fern, mosses and algal plasma membrane ATPase sequences including <i>Arabidopsis thaliana</i> ATPase 1 (UniProt: P20649, TAIR: At2g18960) and ATPase 2 (UniProt: P19456 , TAIR: At4g30190), 3 (UniProt: P20431, TAIR: At5g57350), 4 (UniProt: Q9SU58, TAIR: At3g47950), 6 (UniProt: Q9SH76, TAIR: At2a07560), 7 (UniProt: Q9LY32, TAIR: At3g60330), 8 (UniProt: Q9M2A0, TAIR: At3g42640), 9 (UniProt: Q42556, TAIR: At1g080660), 11 (UniProt: Q9LV11, TAIR: At5g62670) of <i>Arabidopsis thaliana</i> and hydrogen ATPase of <i>Chlamydomonas reinhardtii</i> (Q9FNS3)
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Purity</b>	Antigen affinity purified serum in PBS pH 7.4, conjugated to DyLight® 488.
<b>Format</b>	Liquid
<b>Quantity</b>	40 µg
<b>Storage</b>	Store at 4°C for 12-18 months. A preservative may be added for long time storage up to 2 years.
<b>Additional information</b>	<p>Cellular [compartment marker] for plasma membrane.</p> <p>DyLight® 488 has Amax = 493 nm, Emax = 518 nm. DyLight® is a registered trademark of Thermo Fisher Inc., and its subsidiaries.</p>

### Application information

<b>Recommended dilution</b>	1 : 600-1 : 1000 (IF), 1 : 1000-1 : 2 000 (WB)
<b>Expected   apparent MW</b>	90- 95 kDa ( <i>Arabidopsis thaliana</i> , depending upon an isoform)
<b>Confirmed reactivity</b>	<i>Actinidia chinensis</i> , <i>Aesculus hippocastanum</i> , <i>Arabidopsis thaliana</i> , <i>Camellia sinensis</i> cv. Shu-chazao, <i>Chara australis</i> R.Br, <i>Chlamydomonas reinhardtii</i> , <i>Cucumis sativus</i> , <i>Cucurbita moschata</i> , <i>Glycine max</i> , <i>Kandelia obovata</i> , <i>Hordeum vulgare</i> , <i>Lolium perenne</i> , <i>Lycopersicon esculentum</i> , <i>Malus x domestica</i> Borkh. c.v. Fuji, <i>Marchantia polymorpha</i> , <i>Medicago truncatula</i> , <i>Nicotiana benthamiana</i> , <i>Nicotiana tabacum</i> , <i>Nothaea caeruleascens</i> , <i>Oryza sativa</i> , <i>Petunia hybrida</i> , <i>Phalenopsis Sogo Yukidian cultivar V3</i> , <i>Physcomitrium patens</i> , <i>Picea abies</i> , <i>Pisum sativum</i> , <i>Populus tremula</i> , <i>Pteris vittata</i> (fern), <i>Ricinus communis</i> , <i>Spinacia oleracea</i> , <i>Solanum lycopersicum</i> , <i>Tagetes erecta</i> , <i>Tetraselmis chuii</i> , <i>Zea mays</i> , <i>Vicia faba</i>
<b>Predicted reactivity</b>	Algae, <i>Amaranthus hypochondriacus</i> , <i>Avena sativa</i> , <i>Beta vulgaris</i> , <i>Cyanidioschyzon merolae</i> , <i>Dunaliella spp.</i> , <i>Gossypium hirsutum</i> , <i>Hordeum vulgare</i> , <i>Ostreococcus spp.</i> , <i>Pinus thunbergii</i> , <i>Physcomitrella patens</i> , <i>Mesembrianthemum crystallinum</i> , <i>Mortierella elongata</i> , <i>Nannochloropsis gaditana</i> CCMP526, <i>Ostreococcus tauri</i> , <i>Prosopis alba</i> , <i>Saccharomyces cerevisiae</i> , <i>Solanum tuberosum</i> , <i>Sorghum bicolor</i> , <i>Spinacia oleracea</i> , <i>Triticum aestivum</i> , <i>Ulva prolifera</i> , <i>Ustilago maydis</i>
	Species of your interest not listed? <a href="#">Contact us</a>
<b>Not reactive in</b>	<i>Allium sp.</i> , <i>Aspergillus niger</i> , <i>Citrus limon</i> , <i>Colobanthus apetala</i> , <i>Cuminum cyminum</i> , <i>Curcuma amada</i> , <i>Deschampsia antarctica</i> , <i>Lupinus luteus</i> , <i>Morinda citrifolia</i> , <i>Trigonella foenum</i> , <i>Vicia faba</i>
<b>Additional information</b>	<p><b>VERY IMPORTANT:</b> Please, do not heat up your samples above 70°C as this may cause H<sub>+</sub>ATPase to precipitate, and there will be no signal on your Western Blot.</p> <p>Before SDS-PAGE, centrifuge your samples at room temperature at 10 000 rpm/1 min to remove any aggregates.</p> <p>H<sub>+</sub>ATPase will be less abundant in mature roots and leafs, and therefore detection may require use of very sensitive reagents.</p>
<b>Selected references</b>	To be added when available. Antibody released in October 2023.