

This product is **for research use only** (not for diagnostic or therapeutic use)

contact: support@agrisera.com

Agrisera AB | Box 57 | SE-91112 Vännäs | Sweden | +46 (0)935 33 000 | www.agrisera.com

Product no **AS01 007**

## Anti-Lhca3 | PSI type III chlorophyll a/b-binding protein

### Product information

<b>Immunogen</b>	BSA-conjugated synthetic peptide derived from the Lhca3 protein sequence from <i>Arabidopsis thaliana</i> UniProt: <a href="#">Q9SY97</a> , TAIR: <a href="#">At1g61520</a> . This sequence is highly conserved in Lhcb3 proteins from angiosperms (monocots and dicots) and gymnosperms.
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Purity</b>	Total IgG. Protein G purified in PBS pH 7.4.
<b>Format</b>	Lyophilized
<b>Quantity</b>	0.5 mg
<b>Reconstitution</b>	For reconstitution add 100 µl of sterile water
<b>Storage</b>	Store lyophilized/reconstituted at -20°C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please remember to spin the tubes briefly prior to opening them to avoid any losses that might occur from material adhering to the cap or sides of the tube.

### Application information

<b>Recommended dilution</b>	1 : 2000-1 : 5000 (WB)
<b>Expected   apparent MW</b>	29   25 kDa for <i>Arabidopsis thaliana</i> (due to a transit peptide being cleaved off)
<b>Confirmed reactivity</b>	<i>Arabidopsis thaliana</i> , <i>Aracis hypogae</i> , <i>Bryopsis corticulans</i> , <i>Citrus reticulata</i> , <i>Hordeum vulgare</i> , <i>Oryza sativa</i> , <i>Pisum sativum</i> , <i>Phaseolus vulgaris</i> , <i>Posidonia oceanica</i> , <i>Spinacia oleracea</i> , <i>Triticum aestivum</i> , <i>Triticale</i> , <i>Zea mays</i>
<b>Predicted reactivity</b>	Dicots, Gymnosperms, <i>Glycine max</i>
<b>Not reactive in</b>	No confirmed exceptions from predicted reactivity are currently known
<b>Additional information</b>	Protein is processed into mature form ( <a href="#">Jansson</a> 1999).
<b>Selected references</b>	<p><a href="#">Sarvari</a> et al. (2022). Qualitative and quantitative evaluation of thylakoid complexes separated by Blue Native PAGE. <i>Plant Methods</i>. 2022 Mar 3;18(1):23. doi: 10.1186/s13007-022-00858-2. PMID: 35241118; PMCID: PMC8895881.</p> <p><a href="#">Zhu</a> et al. (2020). A NAC transcription factor and its interaction protein hinder abscisic acid biosynthesis by synergistically repressing NCED5 in <i>Citrus reticulata</i>. <i>J Exp Bot</i>. 2020 Jun 22;71(12):3613-3625.doi: 10.1093/jxb/eraa118.</p> <p><a href="#">Chen</a> et al. (2019). Effects of Stripe Rust Infection on the Levels of Redox Balance and Photosynthetic Capacities in Wheat. <i>Int J Mol Sci</i>. 2019 Dec 31;21(1). pii: E268. doi: 10.3390/ijms21010268.</p> <p><a href="#">Mao</a> et al. (2018). Comparison on Photosynthesis and Antioxidant Defense Systems in Wheat with Different Ploidy Levels and Octoploid Triticale. <i>Int J Mol Sci</i>. 2018 Oct 2;19(10). pii: E3006. doi: 10.3390/ijms19103006.</p> <p><a href="#">Li</a> et al. (2018). Modulating plant growth-metabolism coordination for sustainable agriculture. <i>Nature</i>. 2018 Aug 15. doi: 10.1038/s41586-018-0415-5.</p>