

This product is for research use only (not for diagnostic or therapeutic use)

contact: support@agrisera.com

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | www.agrisera.com

Product no AS15 3085

Anti-LUT5 | beta-carotene hydroxylase

Product information

Immunogen His-tagged, recombinant LUT5 of Arabidopsis thaliana, overexpressed in E.coli, UniProt: Q93VK5, TAIR: AT1G31800

Host Rabbit

Clonality Polyclonal

Purity Serum

Format Lyophilized

Quantity 50 μl

Reconstitution For reconstitution add 50 μl of sterile water

Storage Store lyophilized/reconstituted at -20 °C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please remember to spin the tubes briefly prior to opening them to avoid any losses that might occur from material adhering to

the cap or sides of the tube.

Application information

Recommended dilution 1:2000-1:8000 (WB)

Expected | apparent


66.8 | 64 kDa

Confirmed reactivity | Arabidopsis thaliana

Predicted reactivity | Arabidopsis thaliana

Not reactive in Diatoms

Application example



Total proteins from *Arabidopsis thaliana* leaves, corresponding to 1 µg of chlorophylls of wilde-type (wt) and Lut5 mutant, were extracted with loading buffer (10% glycerol, 62.5 mM Tris pH 6.8, 2% SDS, 5% -mercaptoethanol) and denatured at 100°C (boiling water) for 1 min. Proteins were separated on 15% SDS-PAGE (Laemly) and blotted 1h to PVDF using tank transfer. Blots were blocked with blocking solution (PBS 1X, 0.2% w/v Tween, 5% powder milk) for 1h at room temperature (RT) with agitation. Blot was incubated in the primary antibody diluted in blocking solution, at a dilution of 1: 2,000, 1:4,000, 1:8,000 for 1h at RT with agitation. The antibody solution was decanted and the blot was rinsed briefly twice, then washed 3 times for 10 min in blocking solution at RT with agitation. Blot was incubated in secondary antibody (anti-rabbit IgG alkaline phosphatase conjugated) diluted to 1:30 000 in blocking buffer for 1h at RT with agitation. The blot was washed 2 times for 10 min in blocking solution and once with PBS 1X solution for 10 min, then developed in developing buffer NBT/BCIP by manual agitation.

Courtesy of Stefano Cazzaniga, University of Verona, Italy