## 

This product is for research use only (not for diagnostic or therapeutic use)

contact: support@agrisera.com

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | www.agrisera.com

## Product no AS19 4295 Anti-MS | Malate synthase, (glyoxysomal)

## **Product information**

 Immunogen
 KLH-conjugated peptide derived from Cucurbita maxima UniProt: P24571

 Host
 Rabbit

 Clonality
 Polyclonal

 Purity
 Serum

 Format
 Lyophilized

 Quantity
 50 μl

 Reconstitution
 For reconstitution add 50 μl, of sterile water

 Storage
 Store lyophilized/reconstituted at -20°C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please remember to spin the tubes briefly prior to opening them to avoid any losses that might occur from material adhering to the cap or sides of the tube.

## **Application information**

Recommended dilution	1 : 1000 (WB)
Expected   apparent MW	65 kDa
Confirmed reactivity	Nicotiana tabacum
Predicted reactivity	Arabidopsis thaliana, Cajanus cajan, Cinnamomum micranthum f. kanehirae, Cucumis sativus, Cucurbita maxima, Fagus sylvatica, Glycine max, Jatropha curcas, Morus notabilis, Mucuna pruriens, Parasponia andersonii, Populus alba x tremula, Theobroma cacao, Trema orientale Species of your interest not listed? <u>Contact us</u>
Not reactive in	Sorghum bicolor
Additional information	<b>Experimental contitions:</b> 5 µg of total protein extracted freshly from 3-4 weeks old plant leaves with a blender at 4 °C in 300 mM Sorbitol, 50 mM HEPES, 5mM MgCl <sub>2</sub> . Separated on 10 % SDS-PAGE and blotted 1h to PVDF, semi-dry. Blot was blocked with 6 % milk for 1h 4°C with agitation. Blot was incubated in the primary antibody at a dilution of 1: 1 000 ON at 4°C with agitation.
Selected references	According to <u>South</u> et. al (2019). <u>South</u> et. al (2019). Synthetic glycolate metabolism pathways stimulate crop growth and productivity in the field. Science 2019 Jan 4;363(6422), DOI: 10.1126/science.aat9077