


Product no **AS14 2805****Anti-NBR1 | Autophagy substrate NBR1****Product information**

Immunogen	UBA2 domain of NBR1 of <i>Arabidopsis thaliana</i> , fused with GST, UniProt: Q9SB64 , TAIR: AT4G24690
Host	Rabbit
Clonality	Polyclonal
Purity	Serum
Format	Lyophilized
Quantity	50 µl
Reconstitution	For reconstitution add 50 µl of sterile water
Storage	Store lyophilized/reconstituted at -20°C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please remember to spin the tubes briefly prior to opening them to avoid any losses that might occur from material adhering to the cap or sides of the tube.

Application information

Recommended dilution	1: 1000 (IL), 1 : 500-1 : 5000 (WB)
Expected apparent MW	75 100 kDa
Confirmed reactivity	<i>Arabidopsis thaliana</i> , <i>Physcomitrium patens</i>
Predicted reactivity	<i>Brassicaceae</i> family, <i>Zea mays</i>
	Species of your interest not listed? Contact us
Not reactive in	<i>Citrus sinensis</i> , <i>Nicotiana tabacum</i>
Additional information	Specific extraction method and tissue type needs to be used as described in Minina et al, (2013), Dilution in western blot depends upon amount of NBR1 in the sample
Selected references	<p>Guan et al. (2025). Arabidopsis phospholipase D 2 facilitates vacuolar acidification and autophagy under phosphorus starvation by interacting with VATD. Cell Rep. 2025 Jul 15;44(7):116024. doi: 10.1016/j.celrep.2025.116024.</p> <p>Hoffman-Sommer et al. (2025). The TRAPPC8/TRS85 subunit of the Arabidopsis TRAPPIII tethering complex regulates endoplasmic reticulum function and autophagy. Plant Physiol. 2025 Mar 1;197(3):kiaf042. doi: 10.1093/plphys/kiaf042.</p> <p>Julian et al. (2025). ATG8ylation of vacuolar membrane protects plants against cell wall damage. Nat Plants. 2025 Feb;11(2):321-339. doi: 10.1038/s41477-025-01907-z.</p> <p>Yan et al. (2024). Autophagy modulates Arabidopsis male gametophyte fertility and controls actin organization. Nat Commun. 2024 Nov 21;15(1):10071. doi: 10.1038/s41467-024-54468-8.</p> <p>Mosesso et al. (2024). Arabidopsis CaLB1 undergoes phase separation with the ESCRT protein ALIX and modulates autophagosome maturation. Nat Commun. 2024 Jun 19;15(1):5188. doi: 10.1038/s41467-024-49485-6.</p> <p>Lan et al. (2024). Clathrin Light Chains negatively regulate plant immunity by hijacking the autophagy pathway. Plant Commun. 2024 Apr 30;100937. doi: 10.1016/j.xplc.2024.100937.</p> <p>Rodriguez et al. (2020). Autophagy mediates temporary reprogramming and dedifferentiation in plant somatic cells. bioRxiv doi.org/10.1101/747410</p> <p>Calero-Muñoz et al. (2019). Cadmium induces reactive oxygen species-dependent pexophagy in Arabidopsis leaves. Plant Cell Environ. 2019 Sep;42(9):2696-2714. doi: 10.1111/pce.13597.</p> <p>Jia et al. (2019). Noncanonical ATG8-ABS3 interaction controls senescence in plants. Nat Plants. 2019 Feb;5(2):212-224. doi: 10.1038/s41477-018-0348-x.</p> <p>Hackenberg et al. (2013). Catalase and NO CATALASE ACTIVITY1 promote autophagy-dependent cell death in Arabidopsis. Plant Cell. 2013 Nov;25(11):4616-26. doi: 10.1105/tpc.113.117192. Epub 2013 Nov 27.</p> <p>Minina et al. (2013). Autophagy mediates caloric restriction-induced lifespan extension in Arabidopsis. Aging Cell. 2013 Apr;12(2):327-9. doi: 10.1111/accel.12048. Epub 2013 Feb 28. (method description in supplemental materials)</p>



Total protein from approximately eight 8-d-old seedlings of *Arabidopsis thaliana* was extracted with NuPAGE sample buffer (106 mM Tris-HCl pH 8.5, 141 mM Tris Base, 2% lithium dodecyl sulfate, 0.51 mM EDTA pH 8.0, 10% glycerol, 0.22 mM Coomassie Blue G250, 0.166 mM Phenol Red) with 50 mM dithiothreitol and denatured at 100°C for 5 min. Proteins were separated by electrophoresis on a Bolt 10% Bis-Tris Plus gel (Invitrogen) and transferred for 40 min at 24 V to an Amersham Protran 0.45 µm nitrocellulose membrane (GE Healthcare Life Sciences) using semidry transfer. The membrane was blocked with 8% milk in TBS-T for 2 h at 4°C with agitation. The membrane was incubated in the primary antibody at a dilution of 1:4000 overnight at 4°C with agitation in 8% milk in TBS-T. The antibody solution was decanted and the blot was rinsed briefly twice, then washed 3 times for 5 min in 8% milk in TBS-T at 4°C with agitation. The membrane was incubated in secondary antibody (horse radish peroxidase conjugated goat anti-rabbit IgG) diluted to 1:5000 in 8% milk in TBS-T for 4 h at 4°C with agitation. The membrane was washed 3 times for 5 min in TBS-T at 4°C with agitation and incubated for 2 min with WesternBright chemiluminescent detection reagent (Advansta). Exposure time was 30 seconds.

Courtesy of Pierce Young, Rice University, USA