

This product is **for research use only** (not for diagnostic or therapeutic use)

contact: support@agrisera.com

Agrisera AB | Box 57 | SE-91112 Vännäs | Sweden | +46 (0)935 33 000 | www.agrisera.com

Product no AS16 4064


Anti-NdhB | NAD(P)H-quinone oxidoreductase subunit 2 (chloroplastic)

Product information

Immunogen	KLH-conjugated synthetic peptide derived from <i>Arabidopsis thaliana</i> NdhB protein sequence, UniProt: P0CC32 , P0CC33 , TAIR: AtCg00890 , ATCG01250
Host	Rabbit
Clonality	Polyclonal
Purity	Immunogen affinity purified serum in PBS pH 7.4.
Format	Lyophilized
Quantity	50 µg
Reconstitution	For reconstitution add 50 µl of sterile water
Storage	Store lyophilized/reconstituted at -20°C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please remember to spin the tubes briefly prior to opening them to avoid any losses that might occur from material adhering to the cap or sides of the tube.

Application information

Recommended dilution	1:1000 (WB)
Expected apparent MW	35 kDa
Confirmed reactivity	<i>Arabidopsis thaliana</i> , <i>Hordeum vulgare</i> , <i>Zea mays</i>
Predicted reactivity	<i>Anastatica hierochuntica</i> , <i>Arabis stelleri</i> , <i>Barbarea verna</i> , <i>Braya humilis</i> , <i>Bunias orientalis</i> , <i>Cakile arabica</i> , <i>Camelina sativa</i> , <i>Cannabis sativa</i> , <i>Capsella bursa-pastoris</i> , <i>Capsella rubella</i> , <i>Cardamine impatiens</i> , <i>Chorispora tenella</i> , <i>Cleomella serrulata</i> , <i>Cochlearia pyrenaica</i> , <i>Crucihimalaya wallichii</i> , <i>Dontostemon micranthus</i> , <i>Euclidium syriacum</i> , <i>Eutrema yunnanense</i> , <i>Farselia stylosa</i> , <i>Fragaria ananassa</i> , <i>Hesperis matronalis</i> , <i>Ionopsisidium acaule</i> , <i>Lepidium virginicum</i> , <i>Lobularia maritima</i> , <i>Megadenia pygmaea</i> , <i>Matthiola incana</i> , <i>Nasturtium officinale</i> , <i>Neotorularia korolkowii</i> , <i>Noccaea caerulescens</i> , <i>Olimarabidopsis pumila</i> , <i>Orychophragmus taibaiensis</i> , <i>Oryza sativa</i> , <i>Pachycladon enysii</i> , <i>Phaseolus vulgaris</i> , <i>Physaria ludoviciana</i> , <i>Pugionum cornutum</i> , <i>Schrenkiella parvula</i> , <i>Tarenaya hassleriana</i> , <i>Thlaspi arvense</i> Species of your interest not listed? Contact us
Not reactive in	cyanobacteria, <i>Physcomitrella patens</i>
Selected references	<p>Penzler et al. (2024). A pgr5 suppressor screen uncovers two distinct suppression mechanisms and links cytochrome b6f complex stability to PGR5. <i>Plant Cell</i>. 2024 Mar 27:koae098. doi: 10.1093/plcell/koae098.</p> <p>Penzler et al. (2022) Commonalities and specialties in photosynthetic functions of PROTON GRADIENT REGULATION5 variants in <i>Arabidopsis</i>. <i>Plant Physiol.</i> 2022;190(3):1866-1882. doi:10.1093/plphys/kiac363</p> <p>Shen et al. (2022) Architecture of the chloroplast PSI-NDH supercomplex in <i>Hordeum vulgare</i>. <i>Nature</i>. 2022 Jan;601(7894):649-654. doi: 10.1038/s41586-021-04277-6. Epub 2021 Dec 8. PMID: 34879391.</p> <p>Wada et al. (2021) Identification of a Novel Mutation Exacerbated the PSI Photoinhibition in pgr5/pgrl1 Mutants; Caution for Overestimation of the Phenotypes in <i>Arabidopsis</i> pgr5-1 Mutant. <i>Cells</i>. 2021 Oct 26;10(11):2884. doi: 10.3390/cells10112884. PMID: 34831107; PMCID: PMC8616342.</p>



5µg of total protein from *Zea mays* leaf seedling tissue was extracted with protein extraction buffer (100 mM Tris pH7.5, 10% glycerol, 1 mM EDTA, 1mM EGTA, 2mM pMSF, 0.002 mg leupeptin, 0.002 mg pepstatin A, 2.5 mM DTT) and denatured in 1XPSB (33 mM Tris pH 6.8, 50 mM 2-Mercaptoethanol, 2% SDS, 10% glycerol, 0.1% bromophenol blue) at 70C for 5 min, separated on 4-20% Tris-Glycine SDS-PAGE and blotted overnight to nitrocellulose using tank transfer. Blots were blocked with 10% milk in 1XTBS for 1h at room temperature (RT) with agitation. Blot was incubated in the primary antibody at a dilution of 1:1000 for 1h at RT with agitation in 10% milk in 1XTBS-T. The antibody solution was



This product is **for research use only** (not for diagnostic or therapeutic use)

contact: support@agrisera.com

Agrisera AB | Box 57 | SE-911121 Vännäs | Sweden | +46 (0)935 33 000 | www.agrisera.com

decanted and the blot was rinsed briefly twice, then washed 3 times for 5 min in TBS-T at RT with agitation. Blot was incubated in secondary antibody (anti-rabbit IgG horse radish peroxidase conjugated, from Agrisera, [AS09 602](#)) diluted to 1:10,000 in 10% milk in 1XTBS-T for 1h at RT with agitation. The blot was washed as above and developed for 2 min with chemiluminescent detection reagent.

Courtesy Rosalind Carrier, University of Oregon, USA