

Product no **AS16 4064****Anti-NdhB | NAD(P)H-quinone oxidoreductase subunit 2 (chloroplastic)****Product information**

Immunogen	KLH-conjugated synthetic peptide derived from <i>Arabidopsis thaliana</i> NdhB protein sequence, UniProt: P0CC32 , P0CC33 , TAIR: AtCg00890 , ATCG01250
Host	Rabbit
Clonality	Polyclonal
Purity	Immunogen affinity purified serum in PBS pH 7.4.
Format	Lyophilized
Quantity	50 µg
Reconstitution	For reconstitution add 50 µl of sterile water
Storage	Store lyophilized/reconstituted at -20°C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please remember to spin the tubes briefly prior to opening them to avoid any losses that might occur from material adhering to the cap or sides of the tube.

Application information

Recommended dilution	1:1000 (WB)
Expected apparent MW	35 kDa
Confirmed reactivity	<i>Arabidopsis thaliana</i> , <i>Hordeum vulgare</i> , <i>Zea mays</i>
Predicted reactivity	<i>Anastatica hierochuntica</i> , <i>Arabis stelleri</i> , <i>Barbarea verna</i> , <i>Braya humilis</i> , <i>Bunias orientalis</i> , <i>Cakile arabica</i> , <i>Camelina sativa</i> , <i>Cannabis sativa</i> , <i>Capsella bursa-pastoris</i> , <i>Capsella rubella</i> , <i>Cardamine impatiens</i> , <i>Chorispota tenella</i> , <i>Cleomella serrulata</i> , <i>Cochlearia pyrenaica</i> , <i>Crucihimalaya wallichii</i> , <i>Dontostemon micranthus</i> , <i>Euclidium syriacum</i> , <i>Eutrema yunnanense</i> , <i>Farsetia stylosa</i> , <i>Fragaria ananassa</i> , <i>Hesperis matronalis</i> , <i>Ionopsidium acaule</i> , <i>Lepidium virginicum</i> , <i>Lobularia maritima</i> , <i>Megadenia pygmaea</i> , <i>Matthiola incana</i> , <i>Nasturtium officinale</i> , <i>Neotorularia korolkowii</i> , <i>Noccaea caerulea</i> , <i>Olimarabidopsis pumila</i> , <i>Orychophragmus taibaiensis</i> , <i>Oryza sativa</i> , <i>Pachycladon ensyisii</i> , <i>Phaseolus vulgaris</i> , <i>Physaria ludoviciana</i> , <i>Pugionium cornutum</i> , <i>Schrenkiella parvula</i> , <i>Tarenaya hassleriana</i> , <i>Thlaspi arvense</i> Species of your interest not listed? Contact us
Not reactive in	cyanobacteria, <i>Physcomitrella patens</i>


Selected references

[Penzler](#) et al. (2024). A pgr5 suppressor screen uncovers two distinct suppression mechanisms and links cytochrome b6f complex stability to PGR5. *Plant Cell*. 2024 Mar 27:koae098. doi: 10.1093/plcell/koae098.

[Penzler](#) et al. (2022) Commonalities and specialties in photosynthetic functions of PROTON GRADIENT REGULATION5 variants in Arabidopsis. *Plant Physiol*. 2022;190(3):1866-1882. doi:10.1093/plphys/kiac363

[Shen](#) et al. (2022) Architecture of the chloroplast PSI-NDH supercomplex in *Hordeum vulgare*. *Nature*. 2022 Jan;601(7894):649-654. doi: 10.1038/s41586-021-04277-6. Epub 2021 Dec 8. PMID: 34879391.

[Wada](#) et al. (2021) Identification of a Novel Mutation Exacerbated the PSI Photoinhibition in pgr5/pgl1 Mutants; Caution for Overestimation of the Phenotypes in Arabidopsis pgr5-1 Mutant. *Cells*. 2021 Oct 26;10(11):2884. doi: 10.3390/cells10112884. PMID: 34831107; PMCID: PMC8616342.



5µg of total protein from *Zea mays* leaf seedling tissue was extracted with protein extraction buffer (100 mM Tris pH7.5, 10% glycerol, 1 mM EDTA, 1 mM EGTA, 2mM pMSF, 0.002 mg leupeptin, 0.002 mg pepstatin A, 2.5 mM DTT) and denatured in 1XPSB (33 mM Tris pH 6.8, 50 mM 2-Mercaptoethanol, 2% SDS, 10% glycerol, 0.1% bromophenol blue) at 70C for 5 min, separated on 4-20% Tris-Glycine SDS-PAGE and blotted overnight to nitrocellulose using tank transfer. Blots were blocked with 10% milk in 1XTBS_T for 1h at room temperature (RT) with agitation. Blot was incubated in the primary antibody at a dilution of 1:1000 for 1h at RT with agitation in 10% milk in 1XTBS-T. The antibody solution was

This product is **for research use only** (not for diagnostic or therapeutic use)

contact: support@agrisera.com

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | www.agrisera.com

decanted and the blot was rinsed briefly twice, then washed 3 times for 5 min in TBS-T at RT with agitation. Blot was incubated in secondary antibody (anti-rabbit IgG horse radish peroxidase conjugated, from Agrisera, [AS09_602](#)) diluted to 1:10,000 in 10% milk in 1XTBS-T for 1h at RT with agitation. The blot was washed as above and developed for 2 min with chemiluminescent detection reagent.

Courtesy Rosalind Carrier, University of Oregon, USA