


Product no **AS12 2611****Anti-NRT1,1 | Nitrate transporter 1,1****Product information**

Immunogen	KLH-conjugated synthetic peptide derived from <i>Arabidopsis thaliana</i> NRT1.1 C-terminus, UniProt: Q05085 , TAIR: AT1G12110
Host	Rabbit
Clonality	Polyclonal
Purity	Immunogen affinity purified serum in PBS pH 7.4.
Format	Lyophilized
Quantity	50 µg
Reconstitution	For reconstitution add 50 µl of sterile water
Storage	Store lyophilized/reconstituted at -20°C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please remember to spin the tubes briefly prior to opening them to avoid any losses that might occur from material adhering to the cap or sides of the tube.

Application information

Recommended dilution	1 : 5000 (WB)
Expected apparent MW	64 kDa
Confirmed reactivity	<i>Arabidopsis thaliana</i>
Predicted reactivity	<i>Brassica napus</i> Species of your interest not listed? Contact us
Not reactive in	<i>Alexandrium pacificum</i> , <i>Glycine max</i> , <i>Solanum tuberosum</i>
Selected references	Medici et al. (2015) . AtNIGT1/HRS1 integrates nitrate and phosphate signals at the Arabidopsis root tip. Nat Commun. 2015 Feb 27;6:6274. doi: 10.1038/ncomms7274

application example

50 µg of microsomal protein from wild type *Arabidopsis thaliana* (Col 0) roots at various stages of development or from NRT1.1 deletion mutant (CHL1.5) were solubilized with Laemmli x2 sample buffer and were separated on 10 % SDS-PAGE gel. The proteins were transferred (tank transfer, 100 V, 1h) to a PVDF membrane (Immobilon-P, 0.45 µm, Millipore). Blot was blocked for 2h with agitation at room temperature in blocking buffer (SuperBlock, Thermo) containing 0.05 % Tween 20 (Thermo). Blot was then incubated for 2h at RT and with agitation in the fresh blocking buffer containing the primary antibody at a dilution of 1:5 000. The antibody solution was decanted and the blot was washed 3 times for 10 min in PBS-T (0.05 % Tween 20, Thermo) at RT with agitation. Blot was incubated in secondary antibody (anti-rabbit IgG horse radish peroxidase conjugated, from Agrisera, [AS09 602](#)) diluted to 1:10 000 in blocking buffer, for 2h at RT with agitation. The blot was washed 1 time for 10 min in PBS-T and 2 times for 10 min in PBS. The blot was then incubated with ECL reagent of high sensitivity for 5 min and the chemiluminescence was recorded on LAS 3000, Fuji. Exposure times were between 1-10 min using standard (lowest) sensitivity.

Courtesy of Dr. Wojciech Szponarski, French National Institute of Agricultural Research, France