


Product no **AS22 4879****Anti-OPT3 | Oligopeptide transporter 3****Product information**

Immunogen	KLH-conjugated peptide derived from <i>Arabidopsis thaliana</i> OPT3 protein, UniProt: Q23482 TAIR: AT4G16370
Host	Rabbit
Clonality	Polyclonal
Purity	Antigen affinity purified serum, in PBS pH 7.4
Format	Lyophilized
Quantity	50 µg
Reconstitution	For reconstitution, add 50 µl of sterile or deionized water.
Storage	Store lyophilized/reconstituted at -20°C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please, remember to spin tubes briefly prior to opening them to avoid any losses that might occur from lyophilized material adhering to the cap or sides of the tubes.

Application information

Recommended dilution	1 : 1000 (WB)
Expected apparent MW	82.4 kDa
Confirmed reactivity	<i>Arabidopsis thaliana</i>
Predicted reactivity	<i>Capsella rubella</i> , <i>Camelina sativa</i> For other species: inquire about free sample of AS22 4879 Species of your interest not listed? Contact us
Not reactive in	No confirmed exceptions from predicted reactivity are currently known
Selected references	To be added when available, antibody available in April 2024.



40-60 µg/well of total proteins were extracted from 10-day old *Arabidopsis thaliana* seedlings grown on 0.5X strength Murashige & Skoog media supplemented with 1% sucrose. Col-0 ecotype (wild-type, WT) and opt3-2 T-DNA insertion knock-down mutant ([Khan et al., 2018](#)) were used to test OPT3 protein abundance using Agrisera OPT3 antibody (Cat.No. 224879, Lot.2311). Proteins were denatured at 65°C for 5 min, separated on an 15 % SDS-PAGE and transferred for 70 min at 65V using a tank transfer system to nitrocellulose membrane as described in ([LaMontagne et al., 2016](#)). Blots were blocked with 1X PBS + 0.1% Tween20 (PBS-T) + 5% milk for 2 h at room temperature (RT) with agitation. Primary antibodies were diluted in PBS-T milk to 1: 5,000 and incubated overnight at 4°C with agitation. Primary antibody solutions were decanted, and blots were washed 3 times (7 minutes each) in 1xPBS-T with agitation. Blot was incubated with secondary antibody goat anti-rabbit HRP conjugated ([AS09 602](#), Agrisera) diluted to 1: 7 500 at RT and developed with chemiluminescent detection reagent according to manufacture recommendations. Exposure time was 1 minute on X-ray films. MPK6 antibody served as loading control.

Results: Lower OPT3 band intensity was detected in opt3-2 knock-down mutant compared to *Arabidopsis thaliana* wild-type ecotype Col-0.

Courtesy of Nga Nguyen and Antje Heese (University of Missouri- Columbia, MO, USA)