

This product is for research use only (not for diagnostic or therapeutic use)

contact: support@agrisera.com

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | www.agrisera.com

## Product no AS21 4552

## Anti-PCNA | Proliferating cell nuclear antigen (Saccharomyces cerevisiae)

## **Product information**

Immunogen Recombinant PCNA protein from Saccharomyces cerevisiae, UniProt: P15873

**Host** Rabbit

Clonality Polyclonal

Purity Total immunoglobulin fraction in PBS. Contains 50% glycerol, filter sterilized.

Format Liquid Quantity 100 μg

Storage

Store at -20 °C and avoid temperature below -25 °C; make aliquots to avoid repeated freeze-thaw cycles. Please remember to spin the tubes briefly prior to opening them to avoid any losses that might occur from material adhering to

## **Application information**

Recommended dilution 1: 100 (IP), 1: 1000 (WB)

Expected | apparent 28.9 kDa


Confirmed reactivity Saccharomyces cerevisiae

**Predicted reactivity** Species of your interest not listed? Contact us

Ogiwara et al. (2007). The INO80 chromatin remodeling complex functions in sister chromatid cohesion. Cell Cycle. Selected references

> 2007 May 2;6(9):1090-5. doi: 10.4161/cc.6.9.4130. Epub 2007 May 8. PMID: 17471029. (ChIP) Hishida et al (2006). Functional and physical interaction of yeast Mgs1 with PCNA: impact on RAD6-dependent DNA damage tolerance. Mol Cell Biol. 2006 Jul;26(14):5509-17. doi: 10.1128/MCB.00307-06. PMID: 16809783; PMCID:

> PMC1592726. (Western Blot) lida et al. (2002). PCNA clamp facilitates action of DNA cytosine methyltransferase 1 on hemimethylated DNA. Genes Cells. 2002 Oct;7(10):997-1007. doi: 10.1046/j.1365-2443.2002.00584.x. PMID: 12354094. (Immunoprecipitation)



20 µg of a crude extract of Saccharomyces cerevisiae was separated on a 12.5 % SDS-PAGE and blotted to a PVDF membrane, followed by blotting in 5 % non-fat milk for 1h/RT. Primary antibody was incubated at 1: 1000 1h/RT, followed by washes and incubation with a secondary goat anti-rabbit IgG HRP conjugated antibodies, used at 1: 10 000 1h/RT. Reaction was developed using chemiluminescence following manufacture's recommendations.